

**Original Article****Chemical Properties of Four Valuable Medicinal Plants on Oshtorankoh, Iran**Elham Faryabi<sup>1\*</sup>, Mitra Noori<sup>2</sup>, Amir Mousavi<sup>3</sup> and Ali Ashraf Jafari<sup>4</sup><sup>1</sup> Department of Agriculture, Payam Noor University, Tehran, Iran<sup>2</sup> Department of Biology, Faculty of Science, Arak University, Arak 38156-8-8349-Iran<sup>3</sup> Department of Plant Biotechnology, National Institute of Genetic Engineering and Biotechnology, Tehran, Iran<sup>4</sup> Research Institute of Forests and Rangelands, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran**Article History**

Received: 30 January 2023

Accepted: 15 April 2023

© 2012 Iranian Society of Medicinal Plants.

All rights reserved.

**Keywords**Flavonoids  
Essential oil  
Lamiaceae  
Mint  
Zagros**\*Corresponding author**

E\_Faryabi@Pnu.ac.ir

**ABSTRACT**

Iran has different environmental conditions due to specific geographical. Therefore, exploring essential oil and flavonoids components among different medicinal may be caused discoveries about valuable components. The present study aimed to study flavonoids profile and essential oils composition of four Lamiaceae species i.e. *Stachys lavandulifolia* Vahl, *Teucrium polium* L., *Thymus daenensis* Čelak., and *Ziziphora clinopodioides* Lam. from Oshtorankoh located on Zagros Mountains, Iran. The plants were collected during 2014-2015 and were identified using available references. A chemical study using two-dimensional paper chromatography (2-DPC) and thin-layer chromatography (TLC) showed that the four studied species contained flavone C and C-/O glucosides and flavonoid sulfates. The result of the present study revealed that *T. polium* had high levels of many flavonoids such as apigenin, genistein, isorhamnetin, kaempferol, quercetin, rutin, and vitexin compared to the other studied plants. According to the results of GC/MS analysis, *S. lavandulifolia* was rich in  $\alpha$ -pinene (21.16%) and 1,8-cineol (16.95%), *T. polium* in  $\alpha$ -pinene (12.21%) and sabinene (9.55%), *T. daenensis* in  $\gamma$ -Terpinene (36.98%) and p-Cymene (26.87%), and *Z. clinopodioides* in piperitone (16.84%) and 1,8-cineol (16.11%) components.

**INTRODUCTION**

Mint (Lamiaceae or Labiatae) is a family with about 236 genera and 6900 to 7200 species worldwide, among which about 124 species and subspecies (30%) are endemic to Iran [1] Many of these species are cultivated for their fragrant leaves and attractive flowers. Aerial parts and leaves are the most widely used plant parts. *Nepeta* (76 spp.), *Salvia* (56 spp.), *Stachys* (34 spp.), *Scutellaria* (19 spp.), *Phlomis* (17 spp.), *Eremostachys* (16 spp.), *Thymus* (16 spp.) and *Teucrium* (12 spp.) are the largest genera in Iran [2]. Mint plants are characteristically aromatic, and many of them are cultivated for their essential oils. They are widely used as spices and flavoring in food industries. Their oils are used in the perfumery, cosmetics, and pharmaceutical industries as bactericides, fungicides, virucides, antiparasitics, and pesticides [3]. More than 8000 flavonoid compounds have been identified in plants that afford protection against ultraviolet radiation, pathogens, and herbivores. They are considered health-promoting and disease-preventing dietary

supplements. Epidemiological, clinical, and animal studies reveal that flavonoids may exert protective effects against various disease conditions including cardiovascular disease and cancer. Flavonoids also possess antibacterial, antiviral, and anti-inflammatory effects [4].

A high-level of variation in flavonoids profiles of four *Stachys* species [5]. Anti-inflammatory effects of *Stachys tibetica* polyphenols and flavonoid extracts [6]. The leaves and aerial parts of *Thymus serpyllum* had antidiabetic effects [7]. 55 compounds in *Stachys lavandulifolia* essential oils, of which the major components were  $\alpha$ -pinene (20.1%),  $\beta$ -pinene (12.1%), and spathulenol (7.2%) [8]. Also, myrcene, sabinene,  $\beta$ -phellandrene, and  $\beta$ -caryophyllene were reported as the main components in *S. lavandulifolia* essential oils [9]. Studies on different *S. lavandulifolia* populations indicated that the essential oil compounds can be varied genetically (ecotype), environmentally, and geographically [10]. Chemical compositions and antifungal activity of *T. daenensis*, *T. caramanicus*,

and *Z. clinopodioides* essential oils [11]. Twenty-four components were distinguished in *T. daenensis* essential oil that thymol,  $\alpha$ -terpinene, p-cymene, methyl carvacrol and  $\alpha$ -thujene were the major components [12]. The major essential oil compositions of *T. polium* L. fruits were as follow  $\alpha$ -pinene,  $\beta$ -pinene, limonene, elemol, and cubenol [13]. In addition, eighty-six compounds in *T. polium* essential oil that the main components were  $\alpha$ -pinene,  $\beta$ -pinene, and p-cymene [14]. Antispasmodic effect of aerial part of *T. polium* essential oil on rat isolated ileum in vitro reported [15]. The main chemical compositions of *Z. clinopodioides* aerial part essential oils were reported as carvacrol,  $\alpha$ - &  $\beta$ -pinene)ulegone, cymene, iso-menthone, iso-menthol, menthyl acetate)-menth-3-en-8-ol, neo iso-menthol, limonene, myrcene)iperitenone, 1,8-cineole) iperitone, spathulenol,  $\beta$ -pinene, borneol, germacrene D,  $\gamma$ -terpinene, and  $\beta$ -bourbonene [16]. The antinociceptive effect of *Z. clinopodioides* essential oil and opioidergic system was studied in

male rats [17]. Due to the general tendency to use natural alternative medicines for the prevention and treatment of diseases, the consumption of plants containing flavonoids and essential oils as food and medicine has increased. Therefore, the present study aimed to identification of flavonoids and essential oils compositions of four Lamiaceae species including *Stachys lavandulifolia*, *Teucrium polium*, *T. daenensis*, and *Z. clinopodioides* collected from “Oshtorankoh” protected area in Lorestan Province located on Zagros Mountains, Iran.

## MATERIAL AND METHODS

### Plant Sampling and Preparation

The aerial parts of four Lamiaceae species including *Stachys lavandulifolia*, *T. polium*, *Thymus daenensis*, and *Z. clinopodioides* were collected from “Oshtorankoh” protected area, Lorestan, Iran during 2014-2015 (Table 1).

Herbarium voucher samples of each species were prepared for reference and deposited at the Arak University Herbarium (Fig. 2).

**Table 1** Collection information and aerial parts two-dimensional paper chromatographical data of four studied Lamiaceae.

Voucher samples	Taxon	Altitude (m)	Total flavonoids no.	Flavone C-& C-/O glucosides no.	Flavonoid sulphates no.
CEF02	<i>S. lavandulifolia</i> vahl	2400	6	3	3
CEF55	<i>T. polium</i> L.	1951	6	2	4
CEF70	<i>T. daenensis</i> Celak	2235	9	5	4
CEF73	<i>Z. clinopodioides</i> Lam.	2223	3	2	1

Lorestan with a 28,294 Km<sup>2</sup> area is located in western Iran in the Zagros Mountains (33°58'N, 48°39'E) (Figure 1).



**Fig. 1** The sampling site, Oshtorankoh in Lorestan Province located on Zagros Mountains, Iran.

*T. daenensis* Čelak*Z. clinopodioides* Lam.*S. lavandulifolia* vahl*T. polium* L.

**Fig. 2** Four Lamiaceae species from Oshtorankoh in Lorestan Province located on Zagros Mountains, Iran. The samples were air-dried for isolation and determination of their flavonoids and essential oils.

### Flavonoids

Flavonoids of plants aerial parts were extracted using 70% ethanol and the extracts were vacuum concentrated to dryness in a rotary evaporator. The flavonoids in each extract were isolated and detected using two-dimensional paper and thin-layer chromatography according to Markham (1982) method [18]. Flavonoids extracts were prepared using the acid hydrolysis method adding Chloride acid 2 M hydrochloric acid and placing them in a water bath at 100 °C for 30 min. After cooling, ethyl acetate was added to separate the flavonoid extract from the non-flavonoids extract. The flavonoid extract, soluble in ethyl acetate at 40 °C and 40 m/sec, was distilled off in a vacuum. Co-chromatography with standards was also performed where possible. Flavonoid standards available for comparison during the study were apigenin, chrysin, genistein, hesperidin, isorhamnetin, kaempferol, luteolin, morin, myricetin, naringenin, quercetin, rhamnetin, rutin, tricetin and vitexin (all obtained commercially, Rutin from Merck, apigenin and luteolin from sigma and the rest from Fluka). Cellulose sheet TLC plates were used and after running in three solvents separately, then were viewed in UV245 nm each spot Rf -values and color comparing to standards helped flavonoids identification. Rf values were calculated and in comparison with used flavonoid standards identification and concentration of each spot were determined using Camag UV 254 and 366 nm, chromatographic map, and UV spectroscopy. Also, Perkin-Elmer Lambda 15 UV/Viz Spectrophotometer was used for spots having similar Rf.

### Essential Oil Compositions

A total of 50 g of dried aerial parts were hydro-distilled in a Clevenger type apparatus for 2.5 h in three replications. The EOs were kept at 4 °C in the dark before analysis. The essential oils were injected into a gas chromatograph and the most appropriate temperature planning of the column was determined to complete the separation of essential oils. Then, essential oils were injected into the Gas Chromatograph/Mass Spectrophotometer Aglient 7890 and compound mass spectra were obtained. Essential oils components of each sample were identified using Retention Index (RIs), studying

mass spectra comparing to Reference Spectra Library [19]. In this study, GC including a mass detector Agilent 5975 C, an Electron Ionization source coupled with an Agilent 7890 GC apparatus was used. A 30-meter-long HP-5MS column, 0.25 mm internal diameter, and 0.25  $\mu\text{m}$  internal layer thickness (film) were used. The GC injection site temperature (Inlet) were regulated 280  $^{\circ}\text{C}$ , mass detector ionization source temperature 150  $^{\circ}\text{C}$ , analyzer (Quadrupole) temperature on 230  $^{\circ}\text{C}$ , and the GC/MS intermediate section temperature 280  $^{\circ}\text{C}$  were set. Each extract was inserted into the device through the injection site and the carrier gas (N) was entered into the device column. Then bypassing the sample and gas, extract components are separated and detected by the detector. Results, and graphs were represented and each essential oil constituents' percentages were recounted according to the FID chromatographic peaks areas assuming that all the essential oil compounds comprise 100%.

## RESULTS

### Flavonoids

According to the results, all of the studies samples contained flavonoid sulphates and flavone C- & C-/O-glucosides (Table 2). *T. daenensis* had the most total flavonoids number and *Z. clinopodioides* had the lowest one. Aglycones were not found in any of them. Flavone C- & C-/O-glucosides and flavonoid sulphates were identified flavonoids series in 2-DPC and TLC shows kind of flavonoids. As Table 2 shows rhamnetin, tricetin, and morin were not found in the

four studied species. However, all of them had isorhamnetin, kaempferol, luteolin, orientin, and quercetin. In addition, the samples had apigenin and vitexin (exception for *T. daenensis*). High levels of many flavonoids such as apigenin, genistein, isorhamnetin, kaempferol, quercetin, rutin, and vitexin were observed in *T. polium*.

### Essential Oils

Results of four Lamiaceae taxa aerial parts essential oils studies from Iran using GC/MS showed, *S. lavandulifolia* was rich in  $\alpha$ -Pinene (21.16%) and 1,8-Cineol (16.95%), *T. polium* in  $\alpha$ -Pinene (12.21%) and Sabinene (9.55%), *T. daenensis* in p-Cymene (26.87%) and  $\gamma$ -Terpinene (36.98%), and *Z. clinopodioides* in 1,8-Cineol (16.11%) and piperitone (16.84%). Carvacrol and piperitone components were just found in *T. daenensis* and *Z. clinopodioides*, respectively. Based on the results, the maximum and minimum percentages of essential oil compositions detected by GC/MS belonged to *T. daenensis* (81.18%) and *T. polium* (59.22%), respectively. 1,8-cineol, (E)-2-decenal, limonene,  $\alpha$ -pinene,  $\beta$ -pinene, and (E)-2-undecenal compositions were found in the four studied species. According to the results, 14 components were detected in the essential oils of the four studied species, among which 9, 11, 10, and 9 components were observed in *S. lavandulifolia*, *T. polium*, *T. daenensis* and *Z. clinopodioides*, respectively. All of the studied species had monoterpene, alcohol, aldehyde, and sesquiterpene while aromatic and fatty acid ester were just found in *S. lavandulifolia* and *T. polium*, respectively (Table 4).

**Table 2** Thin Layer Chromatographical data of four studied Lamiaceae taxa aerial parts flavonoids from Iran.

Species	Flavonoids identification														
	Apigenin	Chrysin	Genistein	Isorhamnetin	Kaempferol	Luteolin	Morin	Myricetin	Naringenin	Orientin	Quercetin	Rhamnetin	Rutin	Tricetin	Vitexin
<i>S. lavandulifolia</i>	+	-	-	+	+	±	-	++	-	++	+	-	++	-	++
<i>T. polium</i>	++	+	++	++	++	+	-	-	-	+	++	-	++	-	++
<i>T. daenensis</i>	-	-	-	+	+	+	-	+++	+	++	+	-	+	-	-
<i>Z. clinopodioides</i>	++	+++	++	+	+	+	-	+	-	+	+	-	-	-	+

Concentration of flavonoids:(nonflavonoid), ± (non or a few flavonoid), + (few flavonoid), ++ (middle concentration of flavonoid), +++ (high concentration of flavonoid).



**Table 3** Comparison of essential oils analysis results of four studied Lamiaceae from Iran using GC/MS.

Component	RI <sup>a</sup>	<i>S. lavandulifolia</i>	<i>T. polium</i>	<i>T. daenensis</i>	<i>Z. clinopodioides</i>
$\alpha$ -Pinene	938	21.16 b	12.21	2.03	3.8
(E)-2-Heptenal	965	4.73	5.85	0	0
Sabinene	979	3.58	9.55	0	4.52
$\beta$ -Pinene	985	3.03	7.87	0.45	3.93
p-Cymene	1037	0	0.13	26.87	0
Limonene	1039	2.44	2.27	0.29	17.4
1,8-Cineol	1041	16.95	0.21	2.66	16.11
$\gamma$ -Terpinene	1069	0	0	36.98	0.93
n-Nonanal	1110	4.88	6.73	0	0
Piperitone	1257	0	0	0	16.84
(E)-2-Decenal	1271	5.46	7.07	1.76	1.5
Thymol	1329	0	0.46	3.63	0
Carvacrol	1339	0	0	4.83	0
(E)-2-Undecenal	1374	5.32	6.87	1.68	1.04
Total (%)	-	67.55	59.22	81.18	66.07

<sup>a</sup> RI: Retention indices .

<sup>b</sup> Values are the relative area percent (peak area relative to total peak area in total ion chromatogram).

**Table 4** Essential oils compounds data of four Lamiaceae taxa aerial parts from Iran using GC/MS.

Compounds	<i>S. lavandulifolia</i>	<i>T. polium</i>	<i>T. daenensis</i>	<i>Z. clinopodioides</i>
EO Compounds N.	9	8	6	4
EO Compounds %	54.76	41.64	91.55	90.43
Total EO N.	56	46	27	30
Total EO %	100	99.98	100	99.99
Monoterpene N.	14	10	14	21
Monoterpene %	53.90	37.92	90.95	89.45
Alcohol N.	5	4	2	1
Alcohol %	2.26	2.18	1.95	0.93
Aldehyde N.	23	21	4	7
Aldehyde %	35.61	47.18	9.38	8.63
Sesquiterpene N.	1	1	1	1
Sesquiterpene %	0.86	3.72	0.6	0.98
n-Alkane N.	3	1	0	0
n-Alkane %	1.43	0.47	0	0
Alkane N.	3	0	2	0
Alkane %	1.43	0	0.71	0
Alkene N.	3	7	0	0
Alkene %	5.09	7.19	0	0
Aromatic N.	1	0	0	0
Aromatic %	0.38	0	0	0
Fatty acid N.	1	1	0	0
Fatty acid %	0.47	0.74	0	0
Fatty acid ester N.	0	1	0	0
Fatty acid ester %	0	0.58	0	0
Ketone N.	0	0	1	0
Ketone %	0	0	0.31	0

## DISCUSSION

Species of the *Teucrium* genus have been used for centuries, the plants are used for different purposes such as diabetes, gastrointestinal disorders, rheumatism, inflammations, tuberculosis, diuretic,

antipyretic, tonic, diaphoretic, analgesic, and antihyperlipidemic [20] In the present study, high levels of many flavonoids such as apigenin, genistein, isorhamnetin, kaempferol, quercetin, rutin, and vitexin were observed in *T. polium*.

Therefore, different applications of these plants are maybe related to these components. The plant also had a low level of luteolin, this component was reported in *Teucrium* species [21]. Different components such as luteolin-7-O-rutinoside and luteolin-7-O-glucoside, which having antioxidant activities, were identified in *T. orientale* [22]. Luteolin 3',4',5,7-tetrahydroxyflavone, is a common flavonoid that has been used in Chinese traditional medicine for treating various diseases such as hypertension and inflammatory, it also has a high level of antioxidant activity (Lin et al., 2008). Our results showed from the isolated compounds of *T. polium*, 47.18% were aldehydes. Monoterpenes with 37.92% consist of  $\alpha$ -pinene,  $\beta$ -pinene, and sabinene are a part of species essential oil [23]. Also,  $\alpha$ -Muurolene,  $\alpha$ -Cadinol, and  $\delta$ -Cadinene were isolated from *T. Polium* [24]. *T. polium* extract enhances the anti-angiogenesis effect of tranilast on human umbilical vein endothelial cells [25]. Based on the results, luteolin, isorhamnetin, kempfrol, myricetin, orientin, quercetin, and rutin were found in *Thymus daenensis*. Luteolin, luteolin 7-O-glucoside, and rosmarinic acid were the most evident of phenolic compounds in the species [26]. The results of GC/MS analysis revealed that the essential oil compounds of the plant containing monoterpenes (90.95%),  $\gamma$ -Terpinene (36.98%), and p-Cymene (26.87%). Previous studies reported the following essential oil components in this species,  $\gamma$ -terpinene,  $\alpha$ -pinene, camphene, myricetin, sabinene hydrate, terpinene, isovaleric acid,  $\delta$ -3-carene, thymol, linalool, thymoquinone, and carvacrol [26]. The carvacrol component has therapeutic potential for the prevention and treatment of colon cancer [27]. In addition, sesquiterpenes (60%) consisting  $\beta$ -caryophyllene,  $\alpha$ -humulene and allo-aromadendrene, alcohols (1.95%), aldehydes (47.18%), alkenes (9.38%), alkanes (0.71%), and ketones (0.31%) were reported in *T. daenensis* (Tables 3 & 4). Growth inhibition of Gram-positive bacteria, i.e. *Staphylococcus aureus*, *Micrococcus luteus*, *Enterococcus faecalis*, *Streptococcus pyogenes* was observed using *T. daenensis* methanolic extract [28]. This growth inhibition of the species on different Gram-positive bacteria could be due present of different components in the essential oil of aerial parts of the plant. The plants in *Stachys* species are consumed as herbal tea in the major parts of the world, and due to

their antibacterial, antifungal, anti-inflammatory, and antioxidant capacity, they are widely used as an herbal remedy in alternative medicine [29]. Flavonoids components are known for their antioxidant properties and one of the most important sources for humans is the diet. Flavonoids make up one of the most pervasive groups of plant phenolics. In this study, high levels of some flavonoid components such as luteolin, myricetin, orientin, rutin, and vitexin were observed in aerial parts of *S. lavandulifolia*. There is a strong positive correlation between total phenolic content and antioxidant activity in this species [29]. Due to different flavonoids contents the species can be used potentially as a readily accessible source of natural antioxidants [30]. The flavonoids were also identified in *S. lavandulifolia* methanolic extract [31]. The essential oil extracted from *S. lavandulifolia* subsp. *lavandulifolia* had antimicrobial and antioxidant activities [32]. The GC/MS analysis showed that the essential oil containing monoterpenes (53.9%) and sesquiterpenes (0.86%). In addition, sabinene,  $\beta$ -pinene,  $\beta$ -myrcene, 3- $\delta$ -carene, limonene, terpinolene, allo-Ocimene, verbenone, trans-carveol, nerol, geraniol, caryophyllene, and caryophyllene oxide components were detected in the essential oil of this plant. Two sesquiterpenes components including caryophyllene and caryophyllene oxide in *S. benthamiana* and *S. officinalis*, respectively [33, 34]. Aerial parts including leaves, flowers, and stem of *Z. clinopodioides* are valuable parts of the plant, are used mostly for traditional food and medicine purposes. It is also used in the preparation of an aromatic tea for gastrointestinal disorders and as an aperitive, carminative, and antiseptic [35]. Our results revealed that the plants were rich in apigenin, chrysin, and genistein. The high level of chrysin component was the previous report in this plant, however, some other components such as kaempferol, myricetin, orientin, and quercetin were also observed [36]. The chrysin flavonoid by different properties such as anti-oxidative and anti-inflammatory effects is a potential prophylactic agent in immunopathological and physicochemical injuries [37]. Therefore, it seems that the plant may have health-promoting and disease-preventing in modern lifestyles. A variation in antioxidant and antimicrobial properties of essential oils of *Z. clinopodioides* accessions which were collected

from different regions in Iran [38]. The antioxidant activity of the plant could be due to the high level of 1,8-cineol in its essential oil [39]. The component also has anti-inflammatory properties used for co-medication in inflammatory airway diseases [40]. Based on the results, the plant was rich in piperitone component, which was not observed in the three other studied plants.

Piperitone component (1  $\mu$ l/ml) enhanced the antimicrobial activity of furazolidone and nitrofurantoin against *Salmonella* spp., *E. coli* strains) *roteus* spp., *Enterobacter* spp., *Citrobacter* spp., *Klebsiella* spp., and *Serratia* spp [41]. In the present study such valuable components including carvacrol, 1,8-cineol)-cymene, (E)-2-decenal, (E)-2-heptenal, limonene, n-nonanal,  $\alpha$ -pinene,  $\beta$ -pinene) piperitone, sabinene,  $\gamma$ -terpinene, thymol, and (E)-2-undecenal were detected in the essential oils of *S. lavandulifolia*, *T. polium*, *T. daenensis*, and *Z. clinopodioides*. The essential oil components can be affected by different factors including plant species, plant genetics, soil conditions, altitude, growth phase) part used, and environmental factors. Therefore, exploring phytochemical variations in different medicinal plants species could lead to discoveries about valuable components.

## CONCLUSION

Iran has different environmental and climatic due to specific geographical and climatic conditions. Therefore, exploring essential oil components among different medicinal may be caused to discoveries about valuable components. There is little information about the essential oil components in medicinal plants from Oshtorankoh region. The result of the present study revealed that *T. polium* had high levels of many flavonoids such as apigenin, genistein, isorhamnetin, kaempferol, quercetin, rutin, and vitexin compared to the other studied plants. According to the results of GC/MS analysis, *S. lavandulifolia* was rich in  $\alpha$ -pinene and 1,8-cineol, *T. polium* in  $\alpha$ -pinene, and sabinene, *T. daenensis* in p-Cymene and  $\gamma$ -Terpinene, and *Z. clinopodioides* in 1,8-cineol and piperitone components.

## REFERENCES

1. Akhmedov A., Rog I., Bachar A., Shomurodov H., Nasirov M., Klein T. Higher risk for six endemic and endangered *Lagochilus* species in Central Asia under drying climate. *Perspectives in Plant Ecology, Evolution and Systematics*. 2021; 48:1255-86.
2. Naghibi F., Mosaddegh M., Mohammadi Motamed S., Ghorbani A. Labiatae family in folk medicine in Iran: from ethnobotany to pharmacology. *Iranian journal of pharmaceutical research (IJPR)*. 2005; 2(2):63-79.
3. Bhatt N., Joshi N., Ghai K. Chemical Composition and in-vitro Antimicrobial Activity of Leaf Essential Oils of *Senecio pedunculatus* Edgew. and *Nepeta coerulescens* Maxim. *Journal of Biologically Active Products from Nature*. 2021; 11(1): 1-10.
4. Karak P. Biological activities of flavonoids: an overview. *International Journal of Pharmaceutical Sciences and Research*. 2019; 10(4):1567-1574.
5. Kharazian N., Mohammadi M. Flavonoid Patterns and their Diversity in ten *Stachys* L. (Lamiaceae) Species from Iran. *Progress in Biological Sci*. 2014;4(2): 203-218.
6. Lim H.S., Kim O.S., Kim B.Y., Jeong S.J. Apigenin from *Scutellaria baicalensis* Georgi inhibits neuroinflammation in BV-2 microglia and exerts neuroprotective effect in HT22 hippocampal cells. *Journal of medicinal food*. 2016; 19(11):1032-1040.
7. Alameer M.N., Mushtaq S., Bashir I, Ullah S., Karim M., Rashid M.N. Malik H., Rashid H. Comparative hypoglycemic activity of different fractions of *Thymus serpyllum* L. in alloxan induced diabetic rabbits. *Pakistan J. Pharmaceutical Sci*. 2016; 29(5)1483-8.
8. Feizbaksh A., Tehrani M.S., Rustaiyan S., Masoudi S. Composition of the essential oil of *Stachys lavandulifolia* Vahl. from Iran. *J. Essential Oil Res*. 2003; 15(2):72-73.
9. Shahnama M., Azami S., Mohammad Hosseini M. Characterization of the essential oil and evaluation of antibacterial activity of methanolic extract of *Stachys lavandulifolia* Vahl. *International Journal of Current Microbiology and Applied Sciences*. 2015; 4(3): 275-283.
10. Pirbalouti A.G., Mohammadi M. Phytochemical composition of the essential oil of different populations of *Stachys lavandulifolia* Vahl. *Asian Pacific J. Tropical Biomedicine*. 2013; 3(2):123-128.
11. Khorasany S., Azizi M.H., Barzegar M., Hamidi Esfahani Z. A study on the chemical composition and antifungal activity of essential oil from *Thymus caramanicus*, *Thymus daenensis* and *Ziziphora clinopodioides*. *Nutr Food Sci Res*. 2016; 3(2):35-42.
12. Rustaiee A.R., Khorshidi J., Tabatabaei M.F., Omidbaigi R., Sefidkon F. Essential oil composition of *Thymus daenensis* Celak. during its phenological cycle. *Journal of Essential Oil-Bearing Plants*. 2010; 13(5):556-560.
13. Sabzeghabaie A, Asgarpanah J. Essential oil composition of *Teucrium polium* L. fruits. *Journal of Essential oil Research*. 2016; 28(1):77-80.
14. Cozzani S., Muselli A., Desjobert J.M., Bernardini A.F., Tomi E., Casanova J. Chemical composition of essential oil of *Teucrium polium* subsp. *capitatum* (L.) from Corsica. *Flavour and Fragrance J*. 2005; 20(4):436-441.

15. Sadraei H., Hajhashemi V., Ghannadi A, Mohseni M. Antispasmodic effect of aerial part of *Teucrium polium* L. essential oil on rat isolated ileum in vitro. Medical J. the Islamic Republic of Iran (MJIRI). 2001; 14(4):355–358.
16. Mahboubi M., Heidary Tabar R., Mahdizadeh S. Chemical composition and antifungal activities of *Ziziphora tenuior* and *Z. clinopodioides* essential oils against dermatophytes. Herba Polonica. 2018; 64(2):37-45.
17. Mohammadifard F., Alimohammadi S. Chemical composition and role of opioidergic system in antinociceptive effect of *Ziziphora clinopodioides* essential oil. Basic and Clinical Neuroscience. 2018; 9(5):357-366.
18. Markham K.R. Techniques of Flavonoid Identification. Academic Press, London, 1982.
19. Adams R. Identification of Essential Oil Components by Gas Chromatography/quadrupole Mass Spectroscopy. Allured publishing corporation, 2001.
20. Jarić S., Mitrović Pavlović P. Ethnobotanical features of *Teucrium* species. %1 içinde *Teucrium* Species: Biology and Applications, M. Stanković, Dü., Springer International Publishing, 2020;111–142.
21. Panovska T K, Kulevanova S, Stefova M. In vitro antioxidant activity of some *Teucrium* species (Lamiaceae). Acta pharm. 2005; 55:207–214.
22. Cakir A., Mavi A., Kazaz C., Yildirim A., Kufrevioglu O. Antioxidant activities of the extracts and components of *Teucrium orientale* L. var. orientale. Turkish Journal of Chemistry. 2006; 30(4):483–494.
23. Amiri H. Antioxidant activity of the essential oil and methanolic extract of *Teucrium orientale* (L.) subsp. *taylori* (Boiss.) Rech. F. Iranian journal of pharmaceutical research: IJPR. 2010; 9(4): 417-23.
24. Guetat A., Al-Ghamdi F.A. Analysis of the essential oil of the germander (*Teucrium polium* L.) aerial parts from the northern region of Saudi Arabia. International Journal of Applied Biology and pharmaceutical Technology. 2014; 5(2): 128-135.
25. Sheikhabahaei F., Khazaei M., Nematollahi-Mahani S.N. *Teucrium polium* extract enhances the anti-angiogenesis effect of tranilast on human umbilical vein endothelial cells. Advanced pharmaceutical bulletin. 2018; 8(1)p:131-139.
26. Sonmezdag A.S., Kelebek H., Selli S. Characterization of aroma-active and phenolic profiles of wild thyme (*Thymus serpyllum*) by GC-MS-Olfactometry and LC-ESI-MS/MS. Journal of food science and technology. 2016; 53(4):1957-1965.
27. Fan K., Li X., Cao Y., Qi Y., Li L., Zhang Q., Sun H. Carvacrol inhibits proliferation and induces apoptosis in human colon cancer cells. Anti-cancer drugs. 2015; 26(8):813–823.
28. Mojab F., Poursaeed M., Mehrgan H., Pakdaman S. Antibacterial activity of *Thymus daenensis* methanolic extract. Pakistan journal of pharmaceutical sciences. 2008; 21(3):210-3.
29. Gören A C. Use of *Stachys* species (mountain tea) as herbal tea and food. Records of Natural Products. 2014; 8(2):71-82.
30. Eghdami A., Sadeghi F., Eradatmand Asli D., Houshmandfar A. Antioxidant activity of methanolic and aqueous extract of *Stachys inflata*. 2011.
31. Rahimi Khoigani S., Rajaei A., Goli S.A.H. Evaluation of antioxidant activity, total phenolics, total flavonoids and LC–MS/MS characterisation of phenolic constituents in *Stachys lavandulifolia*. Natural Product Research. 2017; 31(3):355–358.
32. Lin Y, Shi R, Wang X, Shen H M. Luteolin, a flavonoid with potential for cancer prevention and therapy. Current cancer drug targets. 2008; 8(7):634–646.
33. Karami A. Chemical Constituents of *Stachys benthamiana* Boiss. from Southern Zagros, Iran. Analytical Chemistry Letters. 2016; 6(4). 431–434.
34. Giuliani C, Pellegrino R M, Selvaggi R, Tani C, Tirillini B, Bini L M. Secretory structures and essential oil composition in *Stachys officinalis* (L.) Trevisan subsp. *officinalis* (Lamiaceae) from Italy. Natural product research. 2017;31(9). 1006–1013.
35. Şevket A., Hülya D., Elma T., Alma L., Zia-Ul-Haq M., Hadziabulic A., Hilal A. Chemical composition and antioxidant activity *Ziziphora clinopodioides* ecotypes from Turkey. Romanian Biotechnological Letters. 2016; 21(2). 11298–11303.
36. Tian S., Shi Y., Zhou X., Ge L., Upur H. Total polyphenolic (flavonoids) content and antioxidant capacity of different *Ziziphora clinopodioides* Lam. extracts. Pharmacognosy magazine. 2011;7(25):65-8.
37. Zeinali M., Rezaee S.A., Hosseinzadeh H. An overview on immunoregulatory and anti-inflammatory properties of chrysin and flavonoids substances. Biomedicine & Pharmacotherapy. 2017; 92:998-1009.
38. Shahinfar R., Khanzadi S., Hashemi M., Azzadeh M., Boston A. The effect of *Ziziphora clinopodioides* essential oil and nisin on chemical and microbial characteristics of fish burger during refrigerated storage. Iranian Journal of Chemistry and Chemical Engineering (IJCCE).2017; 36(5):65–75.
39. Al-musawi M.T., Ali A.E.H., Humadi A.A., Al-Kaisei B.I. Antioxidant Effects of 1, 8-Cineole Against Long Term DL-Polychlorinated Biphenyls (PCBs) Toxicity in Domestic Hen's Liver.SRP. Sys Rev Pharm. 2020;11(10):1050–1057.
40. Juergens U. Anti-inflammatory properties of the monoterpene 1.8-cineole: current evidence for co-medication in inflammatory airway diseases. Drug Res. 2014; 64(12):638–646.
41. Shahverdi A.R., Rafei H., Fazeli M., Jamalifar H. Enhancement of antimicrobial activity of furazolidone and nitrofurantoin against clinical isolates of Enterobacteriaceae by piperitone. International Journal of Aromatherapy. 2004; 14(2):77–80.