

## Original Article

# Essential Oils of *Nepeta kurdica* and *Nepeta Chionophila* from Iran as Two Sources of Nepetalactone Isomers

Fatemeh Sefidkon\*, Fatemeh Zakerian, Somayeh Fekri Qomi and Mostafa Golipour

Research Institute of Forests and Rangelands, Agricultural Research Education and Extension Organization (AREEO), Tehran, Iran

### Article History

Received: 01 January 2023  
Accepted: 12 September 2023  
© 2012 Iranian Society of Medicinal Plants.  
All rights reserved.

#### Keywords

*Nepeta kurdica*  
*Nepeta chionophila*  
Essential oil  
Nepetalactone

#### \*Corresponding author

sefidkon@rifr-ac.ir,  
sefidkon@gmail.com

### ABSTRACT

This research aimed to identify the volatile compounds of essential oils found in the aerial parts of two *Nepeta* species (*N. kurdica* Hausskn. & Bornm., and *N. chionophila* Boiss. & Hausskn.). This is the first phytochemical investigation of these two *Nepeta* species. The essential oils were obtained by hydro-distillation and analyzed by a combination of capillary gas chromatography (GC-FID) and gas chromatography-mass spectrometry (GC/MS). Nineteen compounds accounting for 90.8% of *N. kurdica* oil were identified, while there were fifteen volatile components corresponding to 95.1% in *N. chionophila* oil. Nepetalactone isomers make up 64.2% of *N. kurdica* and 90.5% of *N. chionophila* oils. The main component of *N. kurdica* oil was 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (57.7%), while 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (71.8%) and 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (15.9%) were the major constituents of *N. chionophila* oil. So, the essential oils of *N. chionophila* (endemic) and *N. kurdica* (native) can be used the same as catnip (*N. cataria*) oil that is used for antimicrobial, repellent and insecticide effect, as a fortifier, a disinfectant and a cure against cold.

## INTRODUCTION

The genus *Nepeta* L. (Lamiaceae) with about 300 species in the Near East, and central and southern Europe and Asia [1], has 67 species in Iran (more than 50% endemics) with the common Persian name of 'Pune-sa' [2].

The aerial parts of some *Nepeta* species as well as their herbal tea are used as medicinal herbs in Iran, traditionally, for example *N. ispanhica*, *N. binaloudensis*, *N. bracteata* and *N. pogonosperma* [3]. The extracts of some species are also used because of their diuretic property and slight bacteriostatic activity, and in ointments to heal skin disorders of eczema type [4]. Some *Nepeta* species are used in folk medicine as diuretic, diaphoretic, antitussive, antispasmodic, antiasthmatic, febrifuge, emmenagogue and sedative agents [5].

The most intensively studied and well known species is *N. cataria* that is found in the Eastern Mediterranean, Southern Asia, Iran and China, and is commonly known as 'Catnip' because of its irresistible action on cats [5]. The essential oil of *N. cataria* is a rich source of nepetalactone isomers. It is estimated that other *Nepeta* species with the same

essential oil composition (containing high amounts of nepetalactone isomers) can have similar medicinal properties with *N. cataria* that is used as a fortifier, a disinfectant and a cure against cold [4]. Antimicrobial, Repellent and insecticide, Larvicidal, Immunomodulatory, Anticonvulsant and myorelaxant, Analgesic and anti-inflammatory, effects on sexual behavior and other medicinal effects were reported for *N. cataria* and other *Nepeta* species containing nepetalactone isomers in their essential oils [6-8].

So we were interested to study the essential oil content and composition of Iranian *Nepeta* species.

Previous studies showed some Iranian *Nepeta* species contain nepetalactones as the main components of their essential oils, such as *N. eremophila* [9], *N. pogonosperma* [10], *N. meyeri* [11], *N. crassifolia* [12-13], *N. racemosa* [14] and *N. persica* [4].

The major components of *N. pogonosperma* were 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (57.6%) and 1,8-cineole (26.4%). The main constituents of the essential oil of *N. meyeri* were 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (53.2%) and 1,8-cineole (29.3%). The main

constituent of *N. crassifolia* oil was 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (92.6%). The main components of *N. racemosa* oil were 4 $\beta$ -7 $\alpha$ -7 $\beta$ -nepetalactone (33.6%), 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (25.6%) and 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (24.4%). Nepetalactone (up to 80%) was also the main compound in the oil of *N. kotschy* var. *persica* [15]. Of course, there is no nepetalactone isomer in the oils of some other *Nepeta* species such as *N. rivularis* [9], *N. macrosiphon* [16], *N. fissa* [17], *N. satureioides* [18], *N. prostrata* [19], *N. ucrainica* [20] and *N. glomerulosa* [21].

In this research we studied the essential oil content and composition of two native *Nepeta* species, named *N. kurdica* Hausskn. & Bornm, and *N. chionophila* Boiss. & Hausskn. from Iran for the first time.

*N. kurdica* that is synonym with *N. wettsteinii* H. Braun [2] is an annual plant that grows in mountainous areas, steep rocky slopes in Irano-Turanian region of Iran. It is distributed in Azarbaijan and Kurdistan provinces [22]. *N. chionophila* is an endemic herbaceous perennial plant that grows in mountainous areas, in snow patch vegetation at an altitude of 2700-3200m in Irano-Turanian region of Iran. From protection position point of view, both of these species are vulnerable [22].

## MATERIALS AND METHODS

### Plant Material

The aerial parts of *N. kurdica* and *N. chionophila* were collected at full flowering stage, from East Azarbaijan province (Tabriz to Urmia, Islamic Island, altitude 1301m) and Chaharmahal and Bakhtiari province (Zard kuh, altitude 2710m), respectively.



**Fig. 1** Image of *N. kurdica* Hausskn. & Bornm



**Fig. 2** Image of *N. chionophila* Boiss. & Hausskn.

The voucher specimens have been deposited in the national herbarium of Iran (TARI) with herbarium number of 431 and 96910, respectively. Images of collected plants can be seen in Figures 1 and 2.

### Isolation Produce

Air-dried aerial parts of the plants (60-80g) were subjected to hydro-distillation for 3h using a Clevenger-type apparatus. The oils separated from water and dried over anhydrous sodium sulfate and stored in sealed vials at low temperature before analysis.

### GC and GC/MS Analysis

GC analyses were performed using a Shimadzu GC-9A gas chromatograph equipped with a DB-5 fused silica column (30 m x 0.25 mm i.d., film thickness 0.25  $\mu$ m). Oven temperature was held at 50  $^{\circ}$ C for 5 min. and then programmed to 240  $^{\circ}$ C at a rate of 3  $^{\circ}$ C/min. Detector (FID) temperature was 265  $^{\circ}$ C and injector temperature was 250  $^{\circ}$ C. Helium was used as carrier gas with a linear velocity of 32 cm/s.

Gas chromatography–mass spectrometry (GC/MS) analysis was conducted with a Varian 3400 GC connected to a mass spectrometer Saturn model, equipped with a DB-5-fused silica capillary column (30 m length, 0.25 mm internal diameter and 0.25  $\mu$ m film thickness); the injection chamber temperature and the transfer line were set at 260 and 270  $^{\circ}$ C, respectively. The carrier gas was helium with a linear velocity of 31.5 cm/s, split ratio of 1 to 60, 1 s of scan time, and mass range of 40–300 a.m.u. The oven temperature adjustment program was the same as the one mentioned for GC.

The percentages of compounds were calculated by the area normalization method, without considering response factors. The retention indices were calculated for all volatile constituents using a homologous series of n-alkanes. The components of

the oil were identified by comparison of their mass spectra with those of a computer library or with authentic compounds and confirmed by comparison of their retention indices either with those of authentic compounds or with data published in the literature [23-25].

## RESULTS AND DISCUSSION

### Plant Materials and Essential Oil Extraction and Analysis

The aerial parts of two Iranian *Nepeta* species named *N. kurdica* Hausskn. & Bornm, and *N. chionophila* Boiss. & Hausskn. were collected from their natural habitats (Azarbaijan and Chaharmahal and Bakhtiari Provinces, respectively) and their

essential oils were obtained by hydro-distillation. Essential oil yields were calculated based on dry weights of plant materials. The oils were analyzed by a combination of capillary gas chromatography, using flame ionization detector (GC-FID) and gas chromatography-mass spectrometry (GC/MS).

### Essential Oil Content and Composition

The essential oil yields of *N. kurdica* and *N. chionophila* were 0.18% and 0.68% (w/w based on dried plants), respectively. Nineteen components were identified in the oil of *N. kurdica* (representing 90.8% of the oil) with 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepeta lactone (57.7%),  $\beta$ -bourbonene (7.1%), 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (4.9%) and spathulenol (4.2%) as main constituents.

**Table 1** Essential oil composition of *N. kurdica* and *N. chinophyla*

Compound	RI (cal.)	RI (Lit.)	<i>N. kurdica</i> (%)	<i>N. chinophyla</i> (%)
1,8-cineole	1032	1026	-	0.4
(Z)- $\beta$ -ocimene	1040	1032	-	0.1
(E)- $\beta$ -ocimene	1052	1044	-	0.2
$\gamma$ -terpinene	1062	1054	-	0.1
cis sabinene hydrate	1072	1065	-	0.1
Linalool	1100	1095	-	0.1
<i>Trans</i> -sabinol	1142	1137	-	0.1
$\delta$ -terpineol	1170	1162	0.2	-
Terpinen-4-ol	1180	1174	0.8	-
$\alpha$ -terpineol	1192	1186	0.4	-
<i>Trans</i> -carveol	1219	1215	0.7	-
Pulegone	1239	1233	2.3	-
Thymol	1294	1289	2.5	-
4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone	1363	1357	57.7	15.9
$\alpha$ -copaene	1380	1374	0.7	-
4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone	1390	1386	4.9	71.8
$\beta$ -bourbonene	1392	1387	7.1	-
4 $\alpha$ -7 $\beta$ -7 $\alpha$ -nepetalactone	1395	1391	1.6	2.8
E- $\beta$ -farnesene	1460	1454	2.0	2.4
$\gamma$ -muurolene	1483	1478	0.4	-
$\gamma$ -himachalene	1486	1481	2.4	-
$\beta$ -bisabolene	1510	1505	0.6	0.7
<i>trans</i> -calamenene	1525	1521	tr	-
$\delta$ -cadinene	1527	1522	-	0.2
Elemol	1553	1548	-	0.1
1-nor-bourbonene	1567	1563	0.2	-
Spathulenol	1581	1577	4.2	0.1
Caryophyllene oxide	1586	1582	2.1	-
Total			90.8	95.1

RI = Retention indices in elution order from DB-1 column

RI (cal.) = Calculated retention indices on the DB-5 column by co-injection of a homologous series of n-alkanes C<sub>8</sub>-C<sub>24</sub>

RI (Lit.) = Literature retention indices (23-25)

tr = less than 0.05%

Fifteen compounds were identified in the oil of *N. chionophila* (representing 95.1% of the oil) with 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (71.8%) and 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (15.9%) as major components. Another nepetalactone isomer, 4 $\alpha$ -7 $\beta$ -7 $\alpha$ -nepetalactone, was found at 1.6% and 2.8% in the oils of *N. kurdica* and *N. chionophila*, respectively.

Chemical composition of the oils can be seen in Table 1. The components are listed in order of their elution on the DB-5 column.

The results showed the oils of *N. kurdica* and *N. chionophila* contained high amount of nepetalactone isomers, but in different percentage of each isomer.

The composition of *Nepeta* phytochemicals as well essential oil can define the medicinal value of it. Nepetalactone isomers as mentioned are group of the important essential oil compounds in *Nepeta* species to which most of the biological properties of the genus attributed.

Catnip (*N. cataria*) and some other *Nepeta* species produced nepetalactone isomers. Nepetalactones are iridoid monoterpenes with a broad range of biological activities that act as defensive compounds in some flowering plants. Nepetalactones protect these plants from herbivorous insects by functioning as insect repellents. They are also produced by many aphids, in which they are sex pheromones [26].

As observed in the present research, 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone (57.7%) in *N. kurdica* oil, was higher than *N. chionophila* (15.9%), but 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone (4.9%) was lower than *N. chionophila* oil (71.8%). Totally, 64.2% of *N. kurdica* essential oil and 90.5% of *N. chionophila* essential oil were nepetalactone isomers.

Also nepetalactone isomers were reported in the essential oils of *N. eremophila* [9], *N. pogonosperma* [10], *N. meyeri* [11], *N. crassifolia* [12-13], *N. racemosa* [14] and *N. persica* [4], but they were not found in *N. schiraziana* [39], *N. glomerulosa* subsp. *carmanica* [32], *N. depauperata*, *N. oxyodonta* [31], *N. ucrainica* ssp. *kopetdaghensis* [35], *N. pungens* [40], *N. macrosiphon* [33], *N. involucrate* [36], *N. bracteata* [37], and *N. crispa* [38].

Previous research on repellent activity of the *N. cataria* oil and the main iridoid compounds (nepetalactone isomers), against the malaria mosquito, the brown ear tick, *Rhipicephalus appendiculatus*, and the red poultry mite,

*Dermanyssus gallinae* showed the nepetalactone isomers have the potential to be used in human and livestock protection against major pathogen vectors [27].

There are many other references about medicinal properties of nepetalactone isomers [6-8, 28-31]. So it can be suggested that the essential oils of *N. kurdica* and *N. chionophila* as two Iranian *Nepeta* species have the same properties and can be used for those purposes.

## CONCLUSION

In this study, for the first time, the essential oil content and composition of two Iranian *Nepeta* species were investigated and compared. Nineteen and fifteen compounds were identified in the oils of *Nepeta kurdica* Hausskn. & Bornm, and *N. chionophila* Boiss. & Hausskn. that approximately constitute 90.8% and 95.1% of the oils, respectively. Nepetalactone isomers, as the main components of some *Nepeta* species oils, were found in both studied essential oils in high amount, but in different percentage. 4 $\alpha$ -7 $\alpha$ -7 $\beta$ -nepetalactone that found in the essential oil of *N. chionophila* at 71.8%, was observed less than 5% in the oil of *N. kurdica*. Another nepetalactone isomer, 4 $\alpha$ -7 $\alpha$ -7 $\alpha$ -nepetalactone was found at 57.7% in the oil of *N. kurdica*, but it was observed at 15.9% in the oil of *N. chionophila*. Totally, more than 90% of the essential oil of *N. chionophila* consisted of three nepetalactone isomers that is remarkable amount. So the essential oils of *N. chionophila* (endemic) and *N. kurdica* (native) can be used the same as catnip (*N. cataria*) oil that is used for antimicrobial, repellent and insecticide effect, as a fortifier, a disinfectant and a cure against cold.

## ACKNOWLEDGMENTS

The authors would like to acknowledge the financial support given by the Research Institute of Forests and Rangelands for this work. We are grateful to Dr. Ziba Jamzad, Mehrdokht Najafpoor Navayi and other botanist in TARI for helping us to identify plant samples. We also thank Dr Mirza and Dr Azimi to help us for GC/MS analysis. We are grateful to Dr Amin Zeraatkar for taking photo from *N. chionophila* in habitat.

## REFERENCES

1. Jamzad Z., Ingrouille M., Simmonds M. Three new species of *Nepeta* (Lamiaceae) from Iran. *Taxon*. 2003; 52: 93-98.
2. Mozaffarian V. A Dictionary of Iranian Plant Name. Farhang Moaser, Tehran, Iran. 2013.
3. Zargari A. Medicinal Plants. Tehran University Publications. Tehran, Iran. 1990; Vol 4: 4th ed.
4. Javidnia K., Miri M., Safavi F., Azarpira A., Shafiee A. Composition of the essential oil of *Nepeta persica* Boiss. from Iran. *Flavour Fragr J*. 2002; 17: 20-22.
5. Formisano C., Rigano D., Senatore F. Chemical constituents and biological activities of *Nepeta* species. *Chem Biodivers*. 2011; 8:1783-1818.
6. Süntar, I., Nabavi, S.M., Barreca, D., Fischer, N. and Effert, T., Pharmacological and chemical features of *Nepeta* L. genus: Its importance as a therapeutic agent. *Phyto Res*. 2017; 5946, 1-14.
7. Nestorović, J., Mišić, D., Šiler, B., Soković, M., Glamočlija, J., Ćirić, A., Maksimović, V. and Grubišić, D., Nepetalactone content in shoot cultures of three endemic *Nepeta* species and the evaluation of their antimicrobial activity. *Fitoterapia*. 2010; 81(6): 621-626.
8. Asgarpanah J, Sarabian S, Ziarati P. Essential oil of *Nepeta* genus (Lamiaceae) from Iran: a review. *J Essent Oil Res*. 2013; 26(1): 1-12.
9. Sefidkon F., Jamzad Z., Mirza M. Chemical composition of the essential oil of five Iranian *Nepeta* species (*N. crispa*, *N. mahanensis*, *N. ispanhanica*, *N. eremophila* and *N. rivularis*). *Flavour Fragr J*. 2006; 21: 764-767.
10. Sefidkon F., Akbarinia A. Essential oil composition of *Nepeta pogonosperma* Jamzad & Assadi from Iran. *J Essent Oil Res*. 2003; 15: 327-328.
11. Sefidkon F., Shaabani A. Essential oil composition of *Nepeta meyeri* Benth. from Iran, *Flavour Fragr J*. 2004; 19: 236-236.
12. Dabiri M., Sefidkon F. Chemical composition of *Nepeta crassifolia* Boiss. & Buhse oil from Iran. *Flavour Fragr J*. 2003a; 18: 225-227.
13. Matloubi Moghadam F., Hosseini M. Composition of the essential oil from *Nepeta crassifolia* Boiss. & Buhse. *Flavour Fragr J*. 1996; 11: 113-115.
14. Dabiri M., Sefidkon F. Chemical composition of the essential oil of *Nepeta racemosa* Lam. from Iran. *Flavour Fragr J*. 2003b; 18: 157-158.
15. Hadi N., Sefidkon F., Shojaeiyan A., Jafari A.A. Essential oil diversity of 21 populations from Iranian endemic species *Nepeta kotschy* Boiss. *Iran J Med Arom Plants*. 2016; 32(2): 189-202.
16. Ghannadi A., Aghazari F., Mehrabani M., Mohagheghzadeh A., Mehregan I. Quantity and composition of the SDE prepared essential oil of *Nepeta macrosiphon* Boiss., *Iran J Pharma Res*. 2003; 103-105.
17. Sefidkon F., Dabiri M., Alamshahi A. Analysis of the essential oil of *Nepeta fissa* C. A. May from Iran. *Flavour Fragr J*. 2001; 17: 89-90.
18. Hadian J., Sonboli A., Nejad Ebrahimi S., Mirjalili, M.H., Essential oil composition of *Nepeta satureioides* from Iran. *Chem Nat Compd*. 2006; 42: 175-177.
19. Sajjadi S.E., Ghassemi N., Marini M. Chemical Constituents of the Essential Oil of *Nepeta prostrata* from Iran. *J Essen Oil, Bearing Plants*. 2012; 15(4): 557-560.
20. Shakeri A., Khakdan F., Soheili V., Sahebkar A., Rassam G., Asili J. Chemical composition, antibacterial activity, and cytotoxicity of essential oil from *Nepeta ucrainica* L. spp. *Kopetdaghensis*. *Ind Crops Prod*. 2014; 58: 315-321.
21. Sefidkon F. Essential oil composition of *Nepeta glomerulosa* Boiss. from Iran. *J. Essent Oil Res*. 2001; 13: 422-423.
22. Jalili A., Jamzad Z. Red Data Book of Iran. Research Institute of Forests and Rangelands, Tehran, Iran. 1999.
23. Adams R.P. Identification of Essential Oil Components by Gas Chromatography/Quadrupole Mass Spectroscopy, Allurd Publishing Corp. Carol Stream, USA. 2017.
24. Shibamoto T. Retention Indices in Essential Oil Analysis, In: Capillary Gas Chromatography in Essential Oil Analysis. Edits., P. Sandra and C. Bicchi, Huethig, Verlag, New York. 1987.
25. Davies B.W. Gas chromatographic retention indices of monoterpenes and sesquiterpenes on methyl silicone and Carbowax 20M phases. *J Chromatogr*. 1990; 503: 1-24.
26. Lichman B.R., Godden G.T., Hamilton J.P., Palmer L., Kamileen M.O., Zhao D., Vaillancourt B., Wood J.C., Sun M., Kinser T.J., Henry L.K., Lopez C.R., Dudareva N., Soltis D.E., Soltis P.S., Buell C.R., O'Connor S.E. The evolutionary origins of the cat attractant nepetalactone in catnip. *Sci Adv*. 2020; 6(20): eaba0721.
27. Birkett M.A., Hassanali A., Høglund S., Pettersson J., Pickett J.A. Repellent activity of catmint, *Nepeta cataria*, and iridoid nepetalactone isomers against Afro-tropical mosquitoes, ixodid ticks and red poultry mites. *Phytochem*. 2011; 72(1): 109-114.
28. Khan M.A., Jones I., Loza-Reyes E., Cameron M.M., Pickett J.A., Birkett M.A. Interference in foraging behaviour of European and American house dust mites *Dermatophagoides pteronyssinus* and *Dermatophagoides farinae* (Acari: Pyroglyphidae) by catmint, *Nepeta cataria* (Lamiaceae). *Exp Appl Acarol*. 2012; 57(1): 5-74.
29. Zhu J.J., Berkebile D.R., Dunlap C.A., Zhang A., Boxler D., Tangtrakulwanich K., Behle R.W., Baxendale F., Brewer G. Nepetalactones from essential oil of *Nepeta cataria* represent a stable fly feeding and oviposition repellent. *Med Vet Entomol*. 2011; 26(2): 131-8.
30. Reichert W., Ejercito J., Guda T., Dong X., Wu Q., Ray A., Simon J.E. Repellency Assessment of *Nepeta cataria* Essential Oils and Isolated Nepetalactones on *Aedes aegypti*. *Sci Rep*. 2019; 9(1): 1524.

31. Sajjadi S.E., Eskandari B. Chemical constituents of the essential oil of *Nepeta oxyodonta*. Chem Nat Compd. 2005; 41: 175-177.
32. Sajjadi S.E., Ghassemi N. Volatile constituents of *Nepeta glomerulosa* Boiss. subsp. *carmanica*. Flavour Fragr J. 1999; 14: 265-267.
33. Ghannadi A., Aghazari F., Mehrabani M., Mohagheghzadeh A., Mehregan I. Quantity and composition of the SDE prepared essential oil of *Nepeta macrosiphon* Boiss. Iran J Pharm Res. 2003; 2: 103-105.
34. Mehrabani M., Asadipour A., Saber Amoli S. Chemical constituents of the essential oil of *Nepeta depauperata* Benth. from Iran. Daru. 2004; 12: 98-100.
35. Javidnia K, Miri R, Mehregan I, Sadeghpour H. Volatile constituents of the essential oil of *Nepeta ucrainica* L. ssp. *kopetdaghensis* from Iran. Flavour Fragr J. 2005; 20: 219-221.
36. Sonboli A., Salehi P., Allahyari L. Essential oil composition of *Nepeta involucrata* from Iran. Chem Nat Compd. 2005; 41: 683-685.
37. Sefidkon F., Jamzad Z. Essential oil composition of four Iranian *Nepeta* species (*N. cephalotes*, *N. bornmuelleri*, *N. mirzayanii* and *N. bracteata*). J Essent Oil Res. 2007; 19: 262-265.
38. Mojab F., Nickavar B., Hooshdar Tehrani H. Essential oil analysis of *Nepeta crispa* and *N. menthoides* from Iran. Iran J Pharm Sci. 2009; 5: 43-46.
39. Akhgar M.R., Moradalizadeh M. Chemical composition of the essential oils from stems, flowers and leaves of *Nepeta schiraziana* Boiss. Iran J. Med. Arom. Plants. 2012; 28: 28-34.
40. Farjam M.H. Antibacterial activity and composition of essential oil of *Nepeta pungens* Benth. from Iran. J. Appl Pharm Sci. 2012; 02(04): 103-105.