

Analysis of miRNAs and Genes Related to Cardiovascular and Neurological Diseases in *Cicer arietinum* L.

Mehrangiz Ghafari¹, Farzaneh Fazeli², Afsaneh Mirshekari³, Sara Rashki Ghalenoo^{4*}, Ali Reza Mirzaei⁵ and Bahman Fazeli-Nasab^{6*}

¹ Department of Pathology, School of Medicine, Zabol University of Medical Sciences, Zabol, Iran

² Department of Biology, Payame Noor University, Terhran, Iran

³ Department of Pediatrics, School of Medicine, Amir al momenin Hospital, Zabol University of Medical Sciences, Zabol, Iran

⁴ Department of Cardiology, Zabol University of Medical Sciences, Zabol, Iran

⁵ Department of Agronomy and Plant Breeding, Faculty of Agriculture and Natural Resources, University of Mohaghegh Ardabili, Ardabil, Iran

⁶ Department of Agronomy and Plant Breeding, Agriculture Institute, Research Institute of Zabol, Zabol, Iran

Article Info

Article Type

Original Article

Article History

Received: 24 June 2024

Accepted: 02 October 2024

© 2012 Iranian Society of Medicinal Plants.

All rights reserved.

*Corresponding authors

dr.s.rashki@gmail.com

bfazelinasab@gmail.com



ABSTRACT

Chickpea (*Cicer arietinum* L.) is one of the most important economic legumes in the world, with 16 chromosomes, and belongs to the Leguminosae or Fabaceae family. This plant is sensitive to cold and is subjected to many environmental stresses every year. In general, the stress of cold, drought, and salinity, in addition to causing severe damage and yield loss, also leads to changes in physiological processes and gene expression. MicroRNAs are a group of conserved RNAs in plants and animals that play an important role in regulating post-transcriptional factors. Chickpea also plays an effective role in promoting resistance to various diseases such as cardiovascular diseases, diabetes, obesity, and cancer. Therefore, in this study, we investigated micro RNAs effective in resistance to environmental stresses, as well as clinical and bioinformatics investigations in the *MTHFR* and *FOLR1* genes against cardiovascular and neurological diseases. The results of this research showed that various miRNAs such as miR159, miR160, miR166, miR167, miR169, miR171, miR172, miR319, miR393, miR394, and miR396 are effective in creating stress resistance in chickpeas through activating different genes. This research showed that micro RNAs act as inhibitory and modulating factors against various stresses. The studies conducted from the analysis of antibodies in *C. arietinum* showed that the *FOLR1* gene is more active in the extracellular part and the plasma membrane and the *MTHFR* gene in the cytosol compared to other cells in other organs. Also, the analysis of the expression of these genes showed that the *FOLR1* gene is less expressed than the *MTHFR* gene in heart but it is more in the brain. On the other hand, the *MTHFR* gene was more expressed in male tissues, muscle tissues, bone marrow, and lymphoid tissue. Also, GMQE in *FOLR1* (0.79) and *MTHFR* (0.87) genes showed that the three-dimensional structure provides an accurate estimation of these genes activities. Finally, it can be concluded that by conducting studies in the field of tracking miRNAs and effective genes, as well as accurate diagnosis with the help of molecular markers, an effective step can be taken to increase resistance to diseases and stresses and significantly improve the performance of the product.

Keywords: Abiotic stress, Clinical analysis, Disease resistance, *FOLR1* gene, Gene expression, MicroRNAs, *MTHFR* gene

How to cite this paper

Ghafari, M., Fazeli, F., Mirshekari, A., Ghalenoo, S.R., Mirzaei, A.R., Analysis of miRNAs and Genes Related to Cardiovascular and Neurological Diseases in *Cicer arietinum* L. Journal of Medicinal plants and By-Products, 2025; 14(2):177-188. doi: 10.22034/jmpb.2024.366716.1740

INTRODUCTION

Today, chickpea (*Cicer arietinum*) is considered the third-most important legume in the world in terms of consumption [1]. This plant is self-pollinating and belongs to temperate regions. Research has shown that the origin of this plant is related to the southeastern regions of Turkey [2]. In general, and according to the morphological classification, chickpea seeds can be divided into two general types. The first is the desi type, with a thick coating that has small seeds and is usually brown. The second type of cable has a thin cover that has

large grains and is usually seen in a cream or beige color [3, 4].

The legume chickpea is said to be nutrient-dense and contains a range of rich and beneficial ingredients, including carbohydrates, proteins, unsaturated fatty acids, minerals, vitamins, dietary fibers, and various isoflavones [3, 5]. A rich source of both carbs and protein, chickpeas are regarded as having higher-quality protein than other pulses [3]. There are 18 different kinds of amino acids in it, and 8 of them are necessary [5]. Scientific evidence supports the biological activity of chickpeas and their

positive impact on human health as a vital source of nutritious components [6].

Pulses are renowned for their high nutritional content, capacity to improve health, and sustainability—they fix nitrogen in the soil, hence lowering the need for fertilizer. Pulses have a low glycemic index and excellent nutritional content. Their drinks are also a rich source of fiber, minerals, and protein [7].

According to recent research, some legume seed protein extracts, especially the albumins from *Lupinus albus*, can prevent colon cancer cell migration and the metalloproteinase MMP-9, which is linked to inflammation and cancer [8, 9]. (Supplementary file (Table 1, Fig. 1)). Additionally, it has been demonstrated that heat-resistant Pis in soy is more efficient and selective against MMP-9 than non-protein substances (such as isoflavones and saponins) [10]. MMP-9 is a significant biomarker and clinical target for inflammatory and cancerous diseases, and its inhibitors are currently thought to be promising treatment options for a range of human conditions, including cancer, osteoporosis, inflammatory bowel disease (IBD), cardiovascular disease, and even neurological disorders [11, 12]. According to some research, several little investigated legume seeds, such chickpea and lupin, have greater bioactivities than soybean and are highly beneficial in decreasing colitis and cancer spreading in vivo [8, 13].

Chickpea, soybean, pea, and lentil are among the legume sources that include dietary Bowman-Birk inhibitors (BBIs), which have the ability to inhibit or prevent inflammatory and carcinogenic processes in the gastrointestinal system [14, 15]. Additional in vitro and in vivo research demonstrated that phytic acid's antioxidant qualities are helpful in preventing cancer and positively impact the growth suppression of a number of cancer types [16].

Chickpea has a very high nutritional value and contains a huge source of vitamins and minerals such as calcium, magnesium, phosphorus, and potassium, as well as amino acids including lysine, methionine, threonine, valine, and leucine, along with beta-carotene. As we know, legumes are a rich source of folate. Folate, or vitamin B9, is very necessary in the body for the synthesis and reconstruction of DNA, cell division, and the maturation of red blood cells. Therefore, this plant has an effective effect on improving cardiovascular diseases, diabetes, neurological diseases, obesity, and cancer [17-20].

Among the several isoflavones present in chickpea are Biochanin A (BCA), calycosin, formononetin, genistein, trifolirhizin, ononin, and sissotrin. Numerous studies have shown that these compounds have unique biological properties and are essential for treating the clinical problems connected to a variety of diseases, including cancer, hyperlipidemia, osteoporosis, the development of the nervous system, and cardiovascular disorders [21, 22]. Isoflavones are the primary bioactive ingredient

found in sprouted chickpea seeds among these constituents. Their wide range of biological actions, which include antioxidative, estrogenic, insecticidal, piscicidal, antifungal, antibacterial, and contraceptive qualities, have helped them to become extremely important [6]. Chickpea products have demonstrated their ability to significantly lower blood lipoprotein levels and cholesterol accumulation. This has a significant hypolipidemic effect and plays a crucial role in the prevention or treatment of a number of diseases, including cardiovascular disease and atherosclerosis, which are caused by hyperlipidemia [6]. These results suggest that BCA may be a therapeutic option for inflammatory cardiovascular disease [6]. Isoflavones are the primary bioactive elements of sprouted chickpea seeds among these elements. Due to their extensive and varied biological actions, which include antioxidant, estrogenic, insecticidal, piscicidal, antifungal, antibacterial, and contraceptive qualities, they have grown to be quite important [6].

Consuming chickpea products has been demonstrated to help prevent or treat a variety of diseases, including atherosclerosis and cardiovascular disease, which are brought on by high blood lipid levels, as well as cholesterol accumulation and blood lipoprotein levels [23]. A diet high in chickpeas reduces the risk of cardiovascular disease in older people by lowering their levels of total cholesterol (TC) and LDL-C [24].

It has been shown that WCL (wild chickpea lectin) protects DNA in a dose-dependent way. Against a variety of fungal infections, the lectin has demonstrated antifungal action. At a dosage of 10 µg/ml, WCL increased the mitogenic response of mouse spleen cells and inhibited HIV-1 reverse transcriptase at an IC₅₀ of 200 µM. Lectin's anticancer potential has been established against human cancer cell lines. IC₅₀ values of 61.8, 54.4, 37.5, and 44.2 µg/ml have been established for the cell viability assay in HepG2, Ishikawa, MCF-7, and MDA-MB-231 cell lines, respectively [25].

Additionally, a variety of proteins and peptides with antibacterial, antioxidant, tumor growth inhibitory, hypoglycemic, and hypolipidemic activities may be present in raw chickpeas and chickpea processed extracts. Chickpea has been demonstrated to have several biological advantages in addition to having strong hepatoprotective, antihypertensive, antiallergic, anticonvulsant, and positive hemagglutination capabilities [6].

Every year, environmental stress causes severe damage to products, which leads to a significant drop in performance. In addition to causing severe damage and yield loss, cold, drought, and salinity stress also lead to changes in biochemical physiological processes and gene expression. Plants also deal with these stresses through several mechanisms. Factors regulating gene expression and transmission of signals, micro RNAs are among the

mechanisms that ultimately lead to regulating metabolic processes and creating resistance in plants [26-28]. Research has demonstrated that consuming chickpea products can significantly lower blood lipoprotein levels and cholesterol accumulation. This has a hypolipidemic effect and plays a major role in the prevention or treatment of a number of diseases, including cardiovascular disease and atherosclerosis, which are caused by hyperlipidemia [29]. After being produced via the one-carbon-donating metabolism of methionine, the thiol-containing amino acid homocysteine is remethylated to methionine, with folates serving as methyl donors [30]. Reduced cell division, the generation of inflammatory cytokines [31], changed nitric oxide metabolism [32], increased oxidative stress [33], enhanced apoptosis [34], and disrupted methylation processes [35] are among the potential negative consequences of homocysteine buildup and folate shortage.

Genes related to folate-mediated one-carbon metabolism and absorption have been shown to contain many variants. These polymorphisms may affect the flow of folate cofactors between DNA synthesis and methylation processes, which might modify the beneficial effects of folates and other B vitamins involved in the metabolism of methyl groups [29]. The folate methylation cycle, which turns homocysteine into methionine, involves the MTHFR gene. DNA, proteins, and lipids are methylated by the usage of S-adenosylmethionine (SAM), a methyl group donor that is derived from methionine [36]. An amino acid alteration at codon Ala222Val causes the MTHFR 677C/T variant, resulting in an unstable enzyme with decreased activity [37]. The buildup of homocysteine [38] and compromised methylation reactions are the outcomes of this mutation. One of the most prevalent repressor mechanisms of tissue-specific genes is DNA methylation. Therefore, this polymorphism's poor methylation may have an impact on the regulation of genes. Through receptors that bind folate and those that are dispersed in a way particular to certain cells and tissues, folate enters the cell. The FR- α and its mouse ortholog Folr1 are examples of such receptors. In developing embryos, Folr1 is expressed in the neural folds, NT, and yolk sac [39].

There are two categories for the roles that miRNAs from medicinal plants play: intra- and cross-kingdom. The primary focus of cross-kingdom studies on therapeutic plant miRNAs is on treating important human diseases [40, 41]. It was discovered that six *Ocimum basilicum* miRNAs, including miR160, 414, and 869.1, modulated 26 target genes relevant to human cancer [42]. Both Homo sapiens miRNA-378 and Aba-miRNA-9497 in *Atropa belladonna* targeted the 3'-untranslated region (3'-UTR) of the mRNA encoding the zinc-finger transcription factor, which is crucial to nervous systems. 691 ZNF [43].

In *Bacopa monnieri*, the TRAF2 gene is a putative target of miRNAs and the upstream signaling factor in the cancer pathway [44]. Within the kingdom, the primary emphasis of miRNA research on medicinal plants is on the metabolic and synthesis pathways of secondary metabolites. miR-4995, miR-5532, and miR-5368 involved in terpenoid production and culture growth conditions in *Picrorhiza kurroa*, an endangered medicinal plant [45]. Target genes for prenylflavonoid production, growth, and development were found in hop (*Humulus lupulus*) miRNAs responding to viral infection [46]. 45 *Aquilegia coerulea* miRNAs that target genes involved in metabolism and stress responses were found using the comparative genomics method [47]. There isn't currently a well-established study system on miRNAs in medicinal plants. With the support of this evidence, the present research was conducted to investigate micro RNAs of resistance to environmental stresses and also to investigate MTHFR (methylenetetrahydrofolate reductase) and FOLR1 (folate receptor- α) genes effective in creating resistance against cardiovascular and neurological diseases.

MATERIALS AND METHODS

First, the NCBI database was used to retrieve the sequences for the FOLR1 (NM_016729.3) and MTHFR (NM_005957.5) genes. These proteins were 656 amino acids long and 257 amino acids long, respectively. The UCSC database was then used to pinpoint where these genes were exactly located. The MBC database was used to calculate the 3D structure of proteins and the Ramachandran plot, while the ProtScale database was used to calculate the molecular weight and isoelectric point of proteins. Cell comparison of MTHFR and FOLR1 genes and the expression of these genes were analyzed by Human Protein Atlas and OMIM database.

RESULTS

Effects of miRNAs

MiRNAs (microRNAs) are small sequences of 22 to 24 nucleotides that lead to many metabolic processes in plants and animals. MiRNAs are involved in the processes of transcription and regulation of post-transcriptional factors; they are also effective in the processes of gene silencing, methylation, translation, and regulation of gene expression [48-50].

In addition, the effect of miRNAs on key regulatory processes such as cellular regulation, biological regulation, and biotic and abiotic stresses such as drought, salinity, heat, cold, oxidative stress, and stress caused by heavy metals in plants and animals has been proven [51-53]. A group of miRNAs is highly protected in plants, especially in medicago legumes and soybeans. This group includes miR159, miR160, miR167, miR169, miR171, miR172, miR393, and miR396. In chickpeas,

this group, together with miR166, miR319, and miR394, plays an effective role in dealing with environmental stresses. It has been shown the protected and effective miRNAs against drought, salt, and cold stresses, along with the mechanisms and genes effective in inducing resistance in chickpea plants (Table 2 and 3) [54-58].

MTHFR Gene

The *MTHFR* gene with accession number NM_005957.5 is located on chromosome 1 (1p36.22). This gene belongs

to the flavoproteins group, which includes 121 genes. Flavoproteins are mostly found in mitochondria and include a nucleic acid derivative of riboflavin. A large number of biological processes include this category of proteins. These proteins frequently perform photosynthesis, DNA repair, and the removal of radicals that cause oxidative stress. The protein produced by this gene converts 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate.

Table 1 Names of miRNAs from medicinal plants that regulate secondary metabolism and cross-kingdom [41]. This table is distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

| Name | Aim Pathway |
|---|--|
| Cross-kingdom | |
| • MIR2911 | COVID-19, influenza A virus, and tumor proliferation |
| • Gas-miR01, and 02 | Anti-inflammatory |
| • MiR414 | Alzheimer's diseases, diabetes, hypoganglionosis, and inflammatory bowel diseases |
| • Oba-miR156f, and 156t | Bile duct carcinoma, lung cancer, and osteoarthritis |
| • Oba-miR160g | Lung cancer, nephronophthisis, and retinitis pigmentosa |
| • Oba-miR482a | Breast cancer, gastric cancer, and ovarian cancer |
| • MiR869.1 | Alzheimer's diseases, cataracts, and diabetes mellitus |
| • Bmn-miR156, 167h, 172d, and 396g | Immune responses |
| • MiR166 | Glioblastoma, papillary thyroid carcinoma, and secretory breast carcinomas |
| • Cac-miR-29c-5p | Breast cancer, and ovarian cancer |
| • Cac-miR-4723-3p | Prostate cancer, and renal cancer |
| • Cac-miR-548d-3p, 5653, 5780d, and 7009-3p | Tumor proliferation, ovarian clear cell adenocarcinoma, breast cancer, and lung cancer |
| • MiR10206, 5059, 5073, 5272, 6135, oba-miR531, and aba-miRNA-9497 | Tumor proliferation, psoriasis, Alzheimer's disease, epilepsy syndromes, immune responses, retinitis pigmentosa, and central nervous system toxicity |
| Secondary metabolism | |
| • MiR5298b, and 8154 | Phenylpropanoid |
| • Smi-miR396b, and miR408 | Salvianolic acid |
| • MiR5298b, and 8154 | Taxol |
| • MiR160b, ath-MIR160b and smi-miR396b | Tanshinone |
| • MiR156 | Sesquiterpene |
| • MiR035, 1168.2, 1438, 156b, 170, 172i, 1858,1873, 2275, 2673a, 2910, 2919, 396b, 408, 5015, 5021, 5658, 828b, 829.1, 8291, f10132-akr, ain-miR1533c, ain-miR156, ain-miR157, cro-miR397a, cro-miR828a, Cs-miR156, mko-miR159b-3p, mko-miR167c-5p, mko-miR168b, mko-miR5082, mko-miR858, mko-miR8610.1, smi-miR12112, smi-miR397, smi-miR396b, and smi-miR408 | Phenolic compounds |
| • MiR_116, 1194, 1276, 15, 1508, 1900, 2141, 2596, 334, 853, 1134, 1533, 160, 164, 167a, 167b, 171, 172, 172d-3p, 2919, 396a, 398f/g, -4995, 5563-x, 5021, 5658, 6435, 838, ain-miR1525, dfr-miR156b, dfr-miR160a, mko-miR156, mko-miR167a, mko-miR396c, mko-miR396g-5p, mko-miR5082, mko-miR827b, mko-miR858, novel-m0022-5p, pmi-miR6300, pmi-miR6173, pmi-miR530, and pmi-Nov_13 | Terpenoid |
| • MiR159, 159a, 166, 171, 172, 2673a, 390, 396, 858, cro-miR160, EY064998, EY082442, EY107691, EY57163, leaf-miR-477, leaf-miR530, root-miR159, root-miR5140, mko-miR159b-3p, mko-miR5082, mko-miR858, mko-miR8610.1, novel miR_218, novel miR_2432, novel miR2642, novel miR_2924, novel miR_457, and novel miR_853 | Esters |
| • MiR2673a, 396, cro-miR160, pso-miR13, pso-miR2161, and pso-miR408 | Alkaloids |
| • MiR156, 5298b, 8154, and novel miR_47 | Saponins |
| • MiR5072, MIR1446-x, and MIR394-y | Quinone |
| • MiR156, 414, 5015b, and 5021 | Essential oil |
| • NovelmiRNA-191, novelmiRNA-23, and novelmiRNA-58 | Triacylglycerols |
| • Pmi-miR396b and pmi-Nov_12 | Green leaf volatile |
| • MIR845-y | Steroid |
| • MiR5021 | Strictosidine |




Fig. 1 miRNAs' multipurpose function in therapeutic plants [41]. This table is distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Table 2 Targets of miRNAs predicted and patterns of their expression in response to cold, drought, and salt stressors

| miRNAs | Stresses | Target gene | Functions |
|--------|----------------------------------|-----------------|---|
| miR159 | Cold, drought, and salt stresses | GAMYB-like-TCP4 | ABA response, NaCl stress response, floral asymmetry and leaf development. Transcription factor TCP4. Transcription factor EREBP-like protein. Transcription factor MYB80. |
| miR160 | Drought and salt stresses | ARF16 | phases of postgermination and seed germination |
| miR166 | Drought and salt stresses | ATHB-15 | Development of the vascular system, leaves, and axillary meristem |
| miR167 | Drought and salt stresses | ABI5 | Growth of the gynoecium and the stamen |
| miR169 | Drought and salt stresses | NF-YA | Plant growth, timing of flowering, and reaction to various biotic stressors |
| miR171 | Drought and salt stresses | NSP2 | abiotic stress response and floral growth |
| miR172 | Drought and salt stresses | RAP2-7 | blooming period, floral organ classification, and cold stress reaction |
| miR319 | Cold stress | GAMYB-like | F-box protein PP2. Transcription factor GAMYB-like. Transcription factor TCP2. |
| miR393 | Cold, drought, and salt stresses | AFB2- AP2 | susceptibility to virulent bacteria AP2-like ethylene-responsive transcription factor protein AUXIN SIGNALING F-BOX 2-like glycine-rich cell wall structural protein 2-like |
| miR394 | Cold stress | MYB98 | Transcription factor MYB98. glycine-rich cell wall structural protein |
| miR396 | Drought and salt stresses | CP29 | leaf and cotyledon development |

As seen in Table 2, a group of miRNAs makes various regulatory chemicals available to the plant through the roots. A group also modulates the level of plant hormones and indirectly helps to control plant pathogens. MiRNAs also stimulate plant growth through ABA signaling and auxin production and are effective in creating resistance to drought, salinity, and cold [59-62].

Table 3 Genes sequence results of *MTHFR*, and *FOLR1*

| Name | <i>MTHFR</i> | <i>FOLR1</i> |
|-----------------------------|-------------------|-------------------|
| Organism | Homo sapiens | Homo sapiens |
| Accession number nucleotide | NM_005957.5 | NM_016729.3 |
| Accession number protein | NP_005948.3 | NP_057941.1 |
| Compartment | Cytosol | Golgi membrane |
| Gene ID | 4524 | 2348 |
| Chromosome | 1 | 11 |
| Cytogenetic location | 1p36.22 | 11q13.4 |
| Chromosome location bp | 11785723-11805964 | 72189709-72196323 |
| nucleotide length | 7018 bp | 930 bp |
| protein length | 656 aa | 257 aa |
| Molecular weight (Da) | 74596.57 | 29819.13 |
| Isoelectric point | 5.22 | 8.30 |
| Total exon | 12 | 4 |



Fig. 2 a) Chromosome 1, the red area is where the *MTHFR* gene is located (1p36.22). b) Chromosome 11, the red area is where the *FOLR1* gene is located (11q13.4).

Also, this protein catalyzes a common substrate for the remethylation of homocysteine to methionine. Deficiency of this gene causes susceptibility to obstructive cardiovascular diseases, neural tube defects, colon cancer, and acute leukemia. In general, a mutation in this gene is associated with methylene tetrahydrofolate reductase deficiency. The number of amino acids, molecular weight, isoelectric point, number of exons, and other characteristics of the *MTHFR* gene are shown in Table 3. Figure 2a shows the location of this gene on the chromosome.

FOLR1 Gene

The *FOLR1* gene, with accession number NM_016729.3, is located on chromosome 11 (11q13.4). The protein encoded by the *FOLR1* gene has a high affinity for folic acid and is attached to the membrane through a solution or a glycosyl-phosphatidylinositol bond. The mutation in this gene is directly related to the destruction of the nerve network and the lack of brain folate transfer. This gene has two promoters that contain several transcriptions start sites and mediate the transfer of 5-methyltetrahydrofolate into cells. The number of amino acids, molecular weight, isoelectric point, number of exons, and other characteristics of the *FOLR1* gene are shown in Table 3. Figure 2b shows the location of this gene on the chromosome.

D Structure of MTHFR and FOLR1 Proteins

First, the sequences of *MTHFR* and *FOLR1* proteins were determined. Then molecular homology modeling using the ExPasy SWISS-MODEL server led to the three-dimensional structure of *MTHFR* and *FOLR1* proteins based on the most similar 1a02 sample (Fig. 3). The estimation of protein quality was determined based on the GMQE scale. The value of GMQE for the *MTHFR* gene was estimated at 0.87 and for the *FOLR1* gene as 0.79. The GMQE values indicate that the 3D structure is an accurate estimate of the *MTHFR* and *FOLR1* genes.

Then the Ramachandran diagram related to *MTHFR* and *FOLR1* proteins was determined to determine the energy level and stability in terms of two angles ϕ and ψ in the proteins. According to Ramachandran's diagram, Ramachandran's energy level based on amino acids in *MTHFR* protein was estimated to be 97.18%, and in *FOLR1* protein 94.12%. Therefore, the proposed model was in a stable state in terms of energy in the three-dimensional structure of these proteins Fig. 4.

Biological Process in MTHFR Gene

The molecular function of the *MTHFR* gene includes catalytic activities, methylenetetrahydrofolate reductase activity, oxidoreductase activity, macromolecular complex binding, and flavin adenine dinucleotide binding. Also, the *MTHFR* gene is effective in various biological processes such as response to hypoxia, cellular

amino acid metabolic process, methionine metabolic process, blood circulation, regulation of histone methylation, response to vitamin B2, tetrahydrofolate interconversion, response to drugs, response to amino acids, S- Adenosylmethionine metabolic process, tetrahydrofolate metabolic process, folic acid metabolic process, homocysteine metabolic process, oxidation-reduction process, response to interleukin-1, and heterochromatin maintenance. In general, this gene is one of the cellular components of the synapse and cytosol, and for this reason, it is very effective in the neural network. The important paralog of this gene is *MTR*.

Biological Process in FOLR1 Gene

The molecular function of the *FOLR1* gene includes folic acid binding, drug binding, methotrexate binding, signaling receptor activity, and folic acid receptor activity. Also, the *FOLR1* gene is effective in various biological processes such as heart looping, entanglement of signals and folic acid, ER to Golgi vesicle-mediated transport, axon regeneration, and response to axon injury, pharyngeal arch artery morphogenesis and anterior neural tube closure. The important paralog of this gene is *FOLR2*.

MTHFR and FOLR1 Gene Expression Analysis

Gene expression analysis showed that *MTHFR* gene expression is relatively lower than that of the *FOLR1* gene in the body. This gene has the highest level of expression in the brain and a low level of expression in other organs. The *FOLR1* gene also has a balanced expression in the whole body; however, it shows more expression in male tissues, muscle tissue, bone marrow, and lymphoid tissue (Figs. 5 and 6).




Fig. 3 A) Three-dimensional structure of *MTHFR* protein. B) Three-dimensional structure of *FOLR1* protein.




Fig. 4 The right side of the MTHFR protein Ramachandran diagram left side of the FOLR1 protein Ramachandran diagram.

Table 4 Localization of the *MTHFR* and *FOLR1* genes

| <i>MTHFR</i> | | <i>FOLR1</i> | |
|-----------------------|------------|-----------------------|------------|
| Compartment | Confidence | Compartment | Confidence |
| Cytosol | 4 | Extracellular | 5 |
| Nucleus | 3 | Plasma membrane | 5 |
| Extracellular | 3 | Golgi apparatus | 4 |
| Plasma membrane | 3 | Endosome | 4 |
| Endoplasmic reticulum | 2 | Endoplasmic reticulum | 4 |
| Peroxisome | 2 | Nucleus | 4 |
| Mitochondrion | 2 | Lysosome | 2 |
| Cytoskeleton | 2 | Cytosol | 2 |
| Golgi apparatus | 1 | Mitochondrion | 2 |
| Lysosome | 1 | Cytoskeleton | 2 |
| Endosome | 1 | Peroxisome | 1 |

miRNAs such as miR159, miR319, miR393, and miR394, which are activated through the response to abscisic acid and whose target gene is MYB, are very effective in creating resistance to drought and cold.

It has investigated the effect of plant growth-promoting rhizobacteria (PGPR) on chickpea miRNAs in desi and kabuli cultivars. Their results showed that PGPR, through a direct effect on miRNAs ultimately leads to the development of resistance, especially in dealing with cold stress [67]. On the other hand, microRNAs affect

Localization of the *MTHFR* and *FOLR1* Genes

Gene localization is based on the UniProtKB and COMPARTMENTS localization databases, as well as ontologies of cellular components visualized by the Gene Ontology Consortium. These data were obtained based on high-performance microscopy and primary sequence predictions (Fig. 7, Table 4).

DISCUSSION

miRNAs generally cause resistance to salt and drought stress through the activation of resistance genes in the abscisic acid pathway. This type of resistance has been reported in plants such as Arabidopsis rice, poplar, cassava, Chinese plum, tomato, etc. [63-66].

Since creating resistance to cold is considered an important goal in chickpeas, and every year cold causes a lot of damage to this plant, it is possible to use miRNAs related to creating cold resistance in this plant.

metabolic and regulatory processes not only directly but also indirectly by affecting other target genes [68]. In general, according to the investigation of miRNAs and resistance genes, we conclude that miRNAs are activated during flowering and seed germination stages against drought and salt stress and through hormonal responses and transcription factors leading to resistance in the plant.

The protein expression level of gene *MTHFR*




Fig. 5 Results of the study of *MTHFR* gene expression

The protein expression level of gene *FOLR1*




Fig. 6 Results of the study of *FOLR1* gene expression

As the results of this research showed, the *MTHFR* gene is related to cardiovascular diseases, and its deficiency causes Homocystinuria. Defects related to the neural network and susceptibility to thromboembolism is also related to the *MTHFR* gene [69-71]. A deficiency of the *MTHFR* gene causes a disturbance in folate metabolism, which ultimately leads to a deficiency of methylene tetrahydrofolate reductase. Severe neurological deterioration in adults that leads to death is caused by this phenotypic disorder [72]. Diseases related to *MTHFR* include homocystinuria caused by a lack of N-methylene

tetrahydrofolate reductase activity, schizophrenia, Alzheimer's, colon cancer, and male infertility caused by misplaced promoter hypermethylation of this gene. The *FOLR1* gene is involved in migrating neural crest cells during heart formation. Also, this gene is involved in the migration of cardiac neural crest cells during the morphogenesis of the outflow tract. One of the most important things to pay attention to in the *FOLR1* gene is the active presence, diversity, and extent of this gene in different cellular components.




Fig. 7 A) Localization of the *MTHFR* gene. B) Localization of the *FOLR1* gene.

These cellular components include the Golgi membrane, endoplasmic reticulum membrane, plasma membrane, an integral part of the plasma membrane, brush border, cell surface, ER to Golgi transport vesicle membrane, basolateral plasma membrane, apical plasma membrane, transport vesicle, clathrin-coated vesicle, an anchored component of the membrane, anchored part of the external side of the plasma membrane, extracellular exosome, extracellular region, nucleus, and endosome [73-78]. On the other hand, as seen, the *FOLR1* gene is expressed in the brain, and *FOLR1* gene deficiency causes neurodegeneration caused by a lack of cerebral folate transport. As we know, NCFTD is an autosomal recessive disorder caused by brain-specific folate deficiency, which is manifested by movement disorders, epilepsy, and leukodystrophy. Recognition and diagnosis of this disorder through the *FOLR1* gene and treatment with folinic acid can reverse the clinical symptoms and improve brain abnormalities and function [79].

The *MTHFR* gene, located on chromosome 1 (1p36.22), is essential for folate metabolism and the remethylation of homocysteine to methionine, processes critical for

DNA synthesis and repair. Variants in this gene, particularly the C677T and A1298C polymorphisms, have been linked to various health risks, including cardiovascular diseases and neural tube defects, due to their impact on enzyme activity and homocysteine levels [80, 81]. The enzyme encoded by *MTHFR* plays a significant role in reducing oxidative stress, which is vital for maintaining cellular integrity. Furthermore, the gene's expression and polymorphisms highlight its relevance in personalized medicine, as understanding these variations can guide therapeutic interventions in affected individuals [82]. Continued research on *MTHFR* is crucial for elucidating its broader implications in metabolic disorders and disease susceptibility.

CONCLUSION

Finally, it can be concluded that by carrying out studies in the field of tracking microRNAs and effector genes as well as accurate examination and diagnosis with the help of molecular markers, an effective step can be taken to increase resistance to diseases and stresses and improve crop yield loss.

REFERENCES

1. Thushan Sanjeeva W.G., Wanasundara J.P.D., Pietrasik Z., Shand P.J. Characterization of chickpea (*Cicer arietinum* L.) flours and application in low-fat pork bologna as a model system. *Food Res Int.* 2010;43(2):617-626.
2. Varshney R.K., Song C., Saxena R.K., Azam S., Yu S., Sharpe A.G., Cannon S., Baek J., Rosen B.D., Tar'an B. Draft genome sequence of chickpea (*Cicer arietinum*) provides a resource for trait improvement. *Nature Biotechnol.* 2013;31(3):240-246.
3. Jukanti A.K., Gaur P.M., Gowda C., Chibbar R.N. Nutritional quality and health benefits of chickpea (*Cicer arietinum* L.): a review. *British J Nutr.* 2012;108(S1):S11-S26.
4. Pathan F., Trimukhe A., Deshmukh R., Annapure U. A peleg modeling of water absorption in cold plasma-treated Chickpea (*Cicer arietinum* L.) cultivars. *Sci Rep.* 2023;13(1):7857.
5. Zia-Ul-Haq M., Iqbal S., Ahmad S., Imran M., Niaz A., Bhangar M.I. Nutritional and compositional study of Desi chickpea (*Cicer arietinum* L.) cultivars grown in Punjab, Pakistan. *Food Chem.* 2007;105(4):1357-1363.
6. Wang J., Li Y., Li A., Liu R.H., Gao X., Li D., Kou X., Xue Z. Nutritional constituent and health benefits of chickpea (*Cicer arietinum* L.): A review. *Food Res Int.* 2021;150:110790.
7. Duarte C.M., Mota J., Assunção R., Martins C., Ribeiro A.C., Lima A., Raymundo A., Nunes M.C., Ferreira R.B., Sousa I. New alternatives to milk from pulses: chickpea and lupin beverages with improved digestibility and potential bioactivities for human health. *Front Nutr.* 2022;9.
8. Lima A.I.G., Mota J., Monteiro S.A.V.S., Ferreira R.M.S.B. Legume seeds and colorectal cancer revisited: Protease inhibitors reduce MMP-9 activity and colon cancer cell migration. *Food Chem.* 2016;197:30-38.
9. Mota J., Direito R., Rocha J., Fernandes J., Sepodes B., Figueira M.E., Raymundo A., Lima A., Ferreira R.B. Lupinus albus protein components inhibit MMP-2 and MMP-9 gelatinolytic activity in vitro and in vivo. *Int J Mol Sci.* 2021;22(24):13286.

10. Lima A., Oliveira J., Saúde F., Mota J., Ferreira R.B. Proteins in soy might have a higher role in cancer prevention than previously expected: soybean protein fractions are more effective MMP-9 inhibitors than non-protein fractions, even in cooked seeds. *Nutrients* 2017;9(3):201.
11. Safavi F., Rostami A. Role of serine proteases in inflammation: Bowman–Birk protease inhibitor (BBI) as a potential therapy for autoimmune diseases. *Expe Mol Pathol.* 2012;93(3):428-433.
12. Swapna B., Fathima K., Muwayyad H., Khanam M., Salma S., Harsha S., Gayas N. A study to assess the associated risk of developing cardiovascular diseases in chronic kidney disease. *Cell Mol Biomed Rep.* 2024;4(3):150-158.
13. Duarte C.M., Mota J., Assunção R., Martins C., Ribeiro A.C., Lima A., Raymundo A., Nunes M.C., Ferreira R.B., Sousa I. New alternatives to milk from pulses: Chickpea and lupin beverages with improved digestibility and potential bioactivities for human health. *Front Nutr.* 2022;9:852907.
14. Clemente A., del Carmen Arques M. Bowman-Birk inhibitors from legumes as colorectal chemopreventive agents. *World J Gastroenterol.* 2014;20(30):10305-10315.
15. Srikanth S., Chen Z. Plant protease inhibitors in therapeutics-focus on cancer therapy. *Front Pharmacol.* 2016;7:470.
16. Kumar A., Singh B., Raigond P., Sahu C., Mishra U.N., Sharma S., Lal M.K. Phytic acid: Blessing in disguise, a prime compound required for both plant and human nutrition. *Food Res Int.* 2021;142:110193.
17. Thudi M., Gaur P.M., Krishnamurthy L., Mir R.R., Kudapa H., Fikre A., Kimurto P., Tripathi S., Soren K.R., Mulwa R. Genomics-assisted breeding for drought tolerance in chickpea. *Funct Plant Biol.* 2014;41(11):1178-1190.
18. Mirzaei A.R., Shakoory-Moghadam V. Bioinformatics analysis and pharmacological effect of *Stevia rebaudiana* in the prevention of type-2 diabetes. *Cell Mol Biomed Rep.* 2022:64-73.
19. Alajaji S.A., El-Adawy T.A. Nutritional composition of chickpea (*Cicer arietinum* L.) as affected by microwave cooking and other traditional cooking methods. *J Food Compos Anal.* 2006;19(8):806-812.
20. Mathew S.E., Shakappa D., Rengel Z. A review of the nutritional and antinutritional constituents of chickpea (*Cicer arietinum*) and its health benefits. *Crop Pasture Sci.* 2022;73(4):401-414.
21. Gupta R.K., Gupta K., Sharma A., Das M., Ansari I.A., Dwivedi P.D. Health risks and benefits of chickpea (*Cicer arietinum*) consumption. *J Agric Food Chem.* 2017;65(1):6-22.
22. Sumanth N., Soumya P., Tabassum A., Mamatha P., Yamini G., Meghamala K., Pravalika G. A study to assess the comorbidities and complications of polycystic ovarian syndrome. *Cell Mol Biomed Rep.* 2023;3(2):107-113.
23. Gopalan R., Gracias D., Madhavan M. Serum lipid and lipoprotein fractions in bengal gram and biochanin a induced alterations in atherosclerosis. *Indian Heart J.* 1991;43(3):185-189.
24. Zahradka P., Wright B., Weighell W., Blewett H., Baldwin A., O K., Guzman R.P., Taylor C.G. Daily non-soy legume consumption reverses vascular impairment due to peripheral artery disease. *Atherosclerosis* 2013;230(2):310-314.
25. Gupta N., Gautam A.K., Bhagyawant S.S. Biochemical characterisation of lectin from wild chickpea (*Cicer reticulatum* L.) with potential inhibitory action against human cancer cells. *J Food Biochem.* 2019;43(2):e12712.
26. Bello M., Tolaba M.P., Aguerre R.J., Suarez C. Modeling water uptake in a cereal grain during soaking. *J Food Engin.* 2010;97(1):95-100.
27. Shittu T.A., Olaniyi M., Oyekanmi A., Okeleye K. Physical and water absorption characteristics of some improved rice varieties. *Food Bioproc Technol.* 2012;5:298-309.
28. Montanuci F.D., Jorge L.M.d.M., Jorge R.M.M. Kinetic, thermodynamic properties, and optimization of barley hydration. *Food Sci Technol.* 2013;33:690-698.
29. Altmäe S., Stavreus-Evers A., Ruiz J.R., Laanpere M., Syvänen T., Yngve A., Salumets A., Nilsson T.K. Variations in folate pathway genes are associated with unexplained female infertility. *Fertil Steril.* 2010;94(1):130-137.
30. Lucock M. Folic Acid: Nutritional Biochemistry, Molecular Biology, and Role in Disease Processes. *Mol Genet Metabol.* 2000;71(1):121-138.
31. Gmyrek G.B., Sozanski R., Jerzak M., Chrobak A., Wickiewicz D., Skupnik A., Sieradzka U., Fortuna W., Gabrys M., Chelmonska-Soyta A. Evaluation of monocyte chemotactic protein-1 levels in peripheral blood of infertile women with endometriosis. *Eur J Obstet Gynecol Reprod Biol.* 2005;122(2):199-205.
32. Thaler C.D., Epel D. Nitric oxide in oocyte maturation, ovulation, fertilization, cleavage and implantation: a little dab'll do ya. *Curr Pharmaceut Des.* 2003;9(5):399-409.
33. Agarwal A., Gupta S., Sharma R.K. Role of oxidative stress in female reproduction. *Reprod Biol Endocrinol.* 2005;3:1-21.
34. Hussein M.R. Apoptosis in the ovary: molecular mechanisms. *Human Reprod Update* 2005;11(2):162-178.
35. Forges T., Monnier-Barbarino P., Alberto J., Gueant-Rodriguez R., Daval J., Gueant J. Impact of folate and homocysteine metabolism on human reproductive health. *Hum Reprod Update* 2007;13(3):225-238.
36. Yamada K., Strahler J.R., Andrews P.C., Matthews R.G. Regulation of human methylenetetrahydrofolate reductase by phosphorylation. *Proc Natl Acad Sci USA* 2005;102(30):10454-10459.
37. Frosst P., Blom H., Milos R., Goyette P., Sheppard C.A., Matthews R., Boers G., Den Heijer M., Kluijtmans L., Van Den Heuve L. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. *Nature Gen.* 1995;10(1):111-113.
38. Harmon D., Woodside J., Yarnell J., McMaster D., Young I., McCrum E., Gey K., Whitehead A., Evans A. The common 'thermolabile' variant of methylene tetrahydrofolate reductase is a major determinant of mild hyperhomocysteinaemia. *QJM: An Int J Med.* 1996;89(8):571-578.
39. López-Escobar B., Wlodarczyk B.J., Caro-Vega J., Lin Y., Finnell R.H., Ybot-González P. The interaction of maternal diabetes with mutations that affect folate metabolism and how they affect the development of neural tube defects in mice. *Dev Dynam.* 2019;248(10):900-917.
40. Li D., Yang J., Yang Y., Liu J., Li H., Li R., Cao C., Shi L., Wu W., He K. A Timely Review of Cross-Kingdom Regulation of Plant-Derived MicroRNAs. *Front Genet.* 2021;12.
41. Sun M., Xu S., Mei Y., Li J., Gu Y., Zhang W., Wang J. MicroRNAs in Medicinal Plants. *Int J Mol Sci.* 2022;23(18):10477.
42. Patel M., Patel S., Mangukia N., Patel S., Mankad A., Pandya H., Rawal R. *Ocimum basilicum* miRNOME revisited: A cross kingdom approach. *Genomics* 2019;111(4):772-785.

43. Avsar B., Zhao Y., Li W., Lukiw W.J. *Atropa belladonna* expresses a microRNA (aba-miRNA-9497) highly homologous to homo sapiens miRNA-378 (hsa-miRNA-378); both miRNAs target the 3'-untranslated region (3'-UTR) of the mRNA encoding the neurologically relevant, zinc-finger transcription factor ZNF-691. *Cell Mol Neurobiol.* 2020;40(1):179-188.
44. Gadhavi H., Patel M., Mangukia N., Shah K., Bhadresha K., Patel S.K., Rawal R.M., Pandya H.A. Transcriptome-wide miRNA identification of *Bacopa monnieri*: a cross-kingdom approach. *Plant Signal Behav.* 2020;15(1):1699265.
45. Vashisht I., Mishra P., Pal T., Chanumolu S., Singh T.R., Chauhan R.S. Mining NGS transcriptomes for miRNAs and dissecting their role in regulating growth, development, and secondary metabolites production in different organs of a medicinal herb, *Picrorhiza kurroa*. *Planta* 2015;241(5):1255-1268.
46. Mishra A.K., Duraisamy G.S., Matoušek J., Radisek S., Javornik B., Jakse J. Identification and characterization of microRNAs in *Humulus lupulus* using high-throughput sequencing and their response to Citrus bark cracking viroid (CBCVd) infection. *BMC Genomics* 2016;17(1) 919.
47. Puzey J.R., Kramer E.M. Identification of conserved *Aquilegia coerulea* microRNAs and their targets. *Gene* 2009;448(1):46-56.
48. Kanwal N., Al Samarrai O.R., Al-Zaidi H.M.H., Mirzaei A.R., Heidari M.J. Comprehensive analysis of microRNA (miRNA) in cancer cells. *Cell Mol Biomed Rep.* 2023;3(2):89-97.
49. Mirzaei A.R., Fazeli F. Bioinformatics analysis of microtubule-associated protein-1 light chain 3 (MAP1LC3A) and (BECN1) genes in autophagy. *Cell Mol Biomed Rep* 2022;2(3):129-137.
50. Sasani S., Rashidi Monfared S., Mirzaei A.R. Identification of some *Echinophora platyloba* miRNAs using computational methods and the effect of these miRNAs in the expression of *TLN2* and *ZNF521* genes in different human body organs. *Cell Mol Biomed Rep.* 2024;4(1):43-53.
51. Sunkar R., Li Y.-F., Jagadeeswaran G. Functions of microRNAs in plant stress responses. *Trends Plant Sci.* 2012;17(4):196-203.
52. Trindade I., Capitão C., Dalmay T., Fevereiro M.P., Santos D.M.d. miR398 and miR408 are up-regulated in response to water deficit in *Medicago truncatula*. *Planta* 2010;231:705-716.
53. Khraiwesh B., Zhu J.-K., Zhu J. Role of miRNAs and siRNAs in biotic and abiotic stress responses of plants. *Biochim Biophys Acta* 2012;1819(2):137-148.
54. Li B., Duan H., Li J., Deng X.W., Yin W., Xia X. Global identification of miRNAs and targets in *Populus euphratica* under salt stress. *Plant Mol Biol.* 2013;81:525-539.
55. Zhou L., Liu Y., Liu Z., Kong D., Duan M., Luo L. Genome-wide identification and analysis of drought-responsive microRNAs in *Oryza sativa*. *J Exp Bot.* 2010;61(15):4157-4168.
56. Zhou Z.S., Zeng H.Q., Liu Z.P., Yang Z.M. Genome-wide identification of *Medicago truncatula* microRNAs and their targets reveals their differential regulation by heavy metal. *Plant Cell Environ.* 2012;35(1):86-99.
57. Zeng F., Qiu B., Wu X., Niu S., Wu F., Zhang G. Glutathione-mediated alleviation of chromium toxicity in rice plants. *Biol Trace Elem Res.* 2012;148:255-263.
58. Pant B.D., Musialak-Lange M., Nuc P., May P., Buhtz A., Kehr J., Walther D., Scheible W.-R.d. Identification of nutrient-responsive *Arabidopsis* and rapeseed microRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. *Plant Physiol.* 2009;150(3):1541-1555.
59. Vacheron J., Desbrosses G., Bouffaud M.-L., Touraine B., Moëgne-Loccoz Y., Muller D., Legendre L., Wisniewski-Dyé F., Prigent-Combaret C. Plant growth-promoting rhizobacteria and root system functioning. *Front Plant Sci.* 2013;4:356.
60. Yang J., Kloepper J.W., Ryu C.-M. Rhizosphere bacteria help plants tolerate abiotic stress. *Trends Plant Sci.* 2009;14(1):1-4.
61. Navarro L., Dunoyer P., Jay F., Arnold B., Dharmasiri N., Estelle M., Voinnet O., Jones J.D. A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* 2006;312(5772):436-439.
62. Timmusk S., Wagner E.G.H. The plant-growth-promoting rhizobacterium *Paenibacillus polymyxa* induces changes in *Arabidopsis thaliana* gene expression: a possible connection between biotic and abiotic stress responses. *Mol Plant-Microbe Interact.* 1999;12(11):951-959.
63. Vannini C., Locatelli F., Bracale M., Magnani E., Marsoni M., Osnato M., Mattana M., Baldoni E., Coraggio I. Overexpression of the rice *Osmby4* gene increases chilling and freezing tolerance of *Arabidopsis thaliana* plants. *Plant J.* 2004;37(1):115-127.
64. Wang W., Shi T., Ni X., Xu Y., Qu S., Gao Z. The role of miR319a and its target gene *TCP4* in the regulation of pistil development in *Prunus mume*. *Genome* 2018;61(1):43-48.
65. Cheng Z., Lei N., Li S., Liao W., Shen J., Peng M. The regulatory effects of MeTCP4 on cold stress tolerance in *Arabidopsis thaliana*: A transcriptome analysis. *Plant Physiol Biochem.* 2019;138:9-16.
66. López-Galiano M.J., García-Robles I., González-Hernández A.I., Camañes G., Vicedo B., Real M.D., Rausell C. Expression of miR159 is altered in tomato plants undergoing drought stress. *Plants* 2019;8(7):201.
67. Jatan R., Chauhan P.S., Lata C. *Pseudomonas putida* modulates the expression of miRNAs and their target genes in response to drought and salt stresses in chickpea (*Cicer arietinum* L.). *Genomics* 2019;111(4):509-519.
68. Lee R.C., Feinbaum R.L., Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell* 1993;75(5):843-854.
69. Deraet E.R., George T.M., Etchevers H.C., Gilbert J.R., Vekemans M., Speer M.C. Human neural tube defects: Developmental biology, epidemiology, and genetics. *Neurotoxicol Teratol.* 2005;27(3):515-524.
70. Seligsohn U., Lubetsky A. Genetic susceptibility to venous thrombosis. *New England J Med.* 2001;344(16):1222-1231.
71. Varga E., Kujovich J. Management of inherited thrombophilia: guide for genetics professionals. *Clin Genet.* 2012;81(1):7-17.
72. Rosenblatt D.S., Lue-Shing H., Arzoumanian A., Low-Nang L., Matiaszuk N. Methylenetetrahydrofolate reductase (MR) deficiency: Thermolability of residual MR activity, methionine synthase activity, and methylcobalamin levels in cultured fibroblasts. *Biochem Med Metabol Biol.* 1992;47(3):221-225.
73. Grant B.D., Donaldson J.G. Pathways and mechanisms of endocytic recycling. *Nature Rev Mol Cell Biol.* 2009;10(9):597-608.
74. Cottam N.P., Ungar D. Retrograde vesicle transport in the Golgi. *Protoplasma* 2012;249:943-955.
75. Kirchhausen T. Three ways to make a vesicle. *Nature Rev Mol Cell Biol.* 2000;1(3):187-198.
76. Appenzeller-Herzog C., Hauri H.-P. The ER-Golgi intermediate compartment (ERGIC): in search of its identity and function. *J Cell Sci.* 2006;119(11):2173-2183.

77. Vickers K.C., Remaley A.T. Lipid-based carriers of microRNAs and intercellular communication. *Curr Opin Lipidol.* 2012;23(2):91.
78. Witwer K.W., Buzás E.I., Bemis L.T., Bora A., Lässer C., Lötvall J., Nolte-'t Hoen E.N., Piper M.G., Sivaraman S., Skog J. Standardization of sample collection, isolation and analysis methods in extracellular vesicle research. *J Extracell Vesicles* 2013;2(1):20360.
79. Steinfeld R., Grapp M., Kraetzner R., Dreha-Kulaczewski S., Helms G., Dechent P., Wevers R., Grosso S., Gärtner J. Folate receptor alpha defect causes cerebral folate transport deficiency: a treatable neurodegenerative disorder associated with disturbed myelin metabolism. *Am J Human Genet.* 2009;85(3):354-363.
80. Rozen R. Molecular Biology of Methylenetetrahydrofolate Reductase (MTHFR): Interrelationships with Folic Acid, Homocysteine and Vascular Disease. In: Robinson, K. (eds) *Homocysteine and Vascular Disease. Developments in Cardiovascular Medicine*, 2000, vol 230. Springer, Dordrecht. https://doi.org/10.1007/978-94-017-1789-2_1681.
81. Wan L., Li Y., Zhang Z., Sun Z., He Y., Li R. Methylenetetrahydrofolate reductase and psychiatric diseases. *Transl Psychiatry* 2018;8(1):242.
82. Undas A., Chojnowski K., Klukowska A., Łętowska M., Mital A., Młynarski W.M., Musiał J., Podolak-Dawidziak M., Szaśiadek M., Treliński J. Determination and interpretation of "MTHFR" gene mutations in gynecology and internal medicine. *Polish Arch Intern Med.* 2019;129(10):728-732.